

On the Zero Divisor Graphs of a Class of Commutative Completely Primary Finite Rings

Walwenda Shadrack Adero¹ and Owino Maurice Oduor²

¹ Maseno University,
Maseno, KENYA
awalwenda@yahoo.com

² University of Kabianga,
Kericho, KENYA
moduor@kabianga.ac.ke

Abstract

Let R be a Completely Primary Finite Ring with a unique maximal ideal $Z(R)$, satisfying $((Z(R))^{n-1} \neq (0)$ and $(Z(R))^n = (0)$. The structures of the units some classes of such rings have been determined. In this paper, we investigate the structures of the zero divisors of R .

Keywords: Finite rings; Zero divisor graphs.

Classification (MSC2010): 05C25, 05C35.

1 Introduction

It is well known, that in a finite ring with 1 , an element is either a zero divisor or a unit. Let R be a finite ring with 1 . An element $u \in R$ is called a zero divisor, if there exists some nonzero element $v \in R$, such that $uv = 0$. The set of zero divisors in R is denoted by $Z(R)$. The zero divisor graphs considered in this paper are due to Anderson and Livingstone in [3], denoted by $\Gamma(R)$. It is conventional, that these graphs present a better description of the structures of the zero divisors. The rest of the notations to be used in the sequel can be obtained from [1,2,4, 5].

Galois rings are completely primary and the importance of completely primary rings in the structure theory of finite rings cannot be underestimated. This is due to the fact that, a commutative finite ring with identity is expressible as a direct sum of completely primary finite rings. The building blocks for the construction of completely primary finite rings are the Galois rings, which are basically the extensions of rings of integers modulo p^n , where p is a prime integer and n is a positive integer. In the next section, we give the construction of the ring under consideration.

2 The Construction

Let R' be the Galois ring of the form $GR(p^{kr}, p^k)$. For each $i = 1, \dots, h$, let

$u_i \in Z(R)$ and U be an h -dimensional R' -module generated by $\{u_1, \dots, u_h\}$ such that $R = R' \oplus U$ is an additive group. On this group, define multiplication by the following relations;

- If $k = 1, 2$, then $pu_i = u_i u_j = u_j u_i = 0, u_i r' = (r')^{\sigma_i} u_i$.
- If $k \geq 3$, then $p^{k-1} u_i = 0, u_i u_j = p^2 \gamma_{ij}, u_i^k = u_i^{k-1} u_j = u_i u_j^{k-1} = 0$;

$u_i r' = (r')^{\sigma_i} u_i$, where $r', \gamma_{ij} \in R', 1 \leq i, j \leq h, p$ is a prime integer, k and r are positive integers and σ_i is the automorphism associated with u_i . Further, let the generators $\{u_i\}$ for U satisfy the additional condition that if $u_i \in U$, then $pu_i = u_i u_j = 0$.

From the given construction, we notice that R is commutative iff σ_i is an identity automorphism. The ring is also completely primary and the subset of the zero divisors $Z(R)$ satisfies the following: $Z(R) = pR' \oplus U$; $(Z(R))^{n-1} = p^{n-1} R'$ and $(Z(R))^n = (0)$.

Due to the differences in their geometrical and structural properties, we present the results of the graph $\Gamma(R)$ for each characteristic of R .



3 Rings of characteristic p

Proposition 3.1 Let $\text{char } R = p$. Then

- i) $|\Gamma(R)| = p^{rh} - 1$
- ii) $\Gamma(R)$ is complete
- iii) $\Gamma(R) = K_{p^{rh}-1}$.
- iv) $\text{diam}(\Gamma(R)) = 1$.
- v) $\text{gr}(\Gamma(R)) = \{\infty \text{ if } r = 1, h = 0, 1 \text{ and } p = 2, 3.$
- vi) The binding number $b(\Gamma(R)) = \infty$.

Proof.

- i) By the above construction, we have that $R' = GF(p^r)$ and $F = R'/pR'$. So $U = F^h$ is an R' -module generated by u_1, \dots, u_h on the additive group $R = R' \oplus U$. It is clear that $Z(R) = R'u_1 \oplus R'u_2 \oplus \dots \oplus R'u_h$ and every non zero element in $Z(R)$ is of the form $(0, r_1, r_2, \dots, r_h)$, for some $r_i \neq 0$. We show that any element not contained in $Z(R)$ is invertible. So let $(r_0, r_1, r_2, \dots, r_h) \notin Z(R)$. Choose an element say $(s_0, s_1, s_2, \dots, s_h) \notin Z(R)$ such that $(r_0, r_1, r_2, \dots, r_h)(s_0, s_1, s_2, \dots, s_h) = (1, 0, 0, \dots, 0)$. This implies that $r_0 s_0 = 1, s_0 = r_0^{-1}$ and $r_0 s_i + r_i s_0 = 0, s_i = -r_i r_0^{-2}$ for $1 < i \leq h$. So, $(r_0, r_1, r_2, \dots, r_h)^{-1} = (r_0^{-1}, -r_1 r_0^{-2}, -r_2 r_0^{-2}, \dots, -r_h r_0^{-2})$. Since $|R| = |R' \oplus U| = p^{(h+1)r}$, $|Z(R)| = p^{rh}$ and $\Gamma(R) = Z(R)^*$, then $|\Gamma(R)| = p^{rh} - 1$ which establishes (i).
- ii) To establish this, note that the product of every pair $(0, r_1, r_2, \dots, r_h), (0, s_1, s_2, \dots, s_h) \in Z(R) - \{0\}$ is equal to zero so that every pair of vertices in $\Gamma(R)$ are adjacent. Hence $\Gamma(R)$ is complete.
- iii) This is a consequence of (ii), since $\Gamma(R)$ is complete having $p^{rh} - 1$ vertices.
- iv) This follows from (ii).
- v) When $r = 1, h = 0$ or 1 and $p = 2$ or 3 , $n = (p^{rh} - 1) \leq 2$. So $\text{gr}(\Gamma(R)) = \infty$. Otherwise, $n = (p^{rh} - 1) > 2$ and since the completeness of the graph implies that $\text{gr}(\Gamma(R)) = 2\text{diam}(\Gamma(R)) + 1$, the result readily follows.
- vi) Since the set S of minimal degree in $V(\Gamma(R))$ is empty, $b(\Gamma(R)) = \infty$.

4. Rings of characteristic p^2

Proposition 4.1 Let $\text{char } R = p^2$. Then the following hold:

- i) $|\Gamma(R)| = p^{(h+1)r} - 1$
- ii) $\Gamma(R)$ is complete.
- iii) $\Gamma(R) = K_{p^{(h+1)r}-1}$
- iv) $\text{diam}(\Gamma(R)) = 1$
- v) $\text{gr}(\Gamma(R)) = \{\infty \text{ if } r = 1, h = 0, \text{ and } p = 2, 3$



{ 3 elsewhere .

vi) The binding number $b(\Gamma(R)) = \infty$.

Proof.

- i) By the construction, $x \in Z(R)^*$ if and only if x is of the form $(0, \overline{r_1}, \overline{r_2}, \dots, \overline{r_h})$. Now let $x = (r_0, \overline{r_1}, \overline{r_2}, \dots, \overline{r_h}) \notin Z(R)$ be an element in R , then x is invertible and indeed x^{-1} is $(s_0, \overline{s_1}, \dots, \overline{s_h})$ such that $s_0 = r_0^{-1}$ and $s_i = -\overline{r_i} r_0^{-2}$ for $1 \leq i \leq h$. Since $|R| = |R'| \cdot |U| = p^{(h+2)r} |Z(R)| = p^{(h+1)r}$ and $\Gamma(R) = Z(R) - \{0\}$, then $|\Gamma(R)| = p^{(h+1)r} - 1$.
- ii) For all $x, y \in Z(R)^*$, $xy = 0$. So $\Gamma(R)$ is clearly complete.
- iii) This follows from (ii).
- iv) It is clear that for all $x, y \in Z(R)^*$, $d(x, y) = 1$. So $Sup\{d(x, y)\} = 1$ for all $x, y \in \Gamma(R)$. Therefore the result is immediate.
- v) When $r = 1, h = 0$ and $p = 2$ or 3 , $\Gamma(R) = K_3$ then $n = (p^{(h+1)r} - 1) \leq 2$ and hence $\Gamma(R)$ has no cycles. So, $gr(\Gamma(R)) = \infty$. Otherwise, for all $r, h \geq 1$ $n = (p^{(h+1)r} - 1) > 2$, so the completeness of $\Gamma(R)$ implies that $gr(\Gamma(R)) = 2diam(\Gamma(R)) + 1 = 3$, since $diam(\Gamma(R)) = 1$.
- vi) Similar to Proposition 3.1.

Proposition 4.2 Let $char R = p$ or p^2 . Then $\Gamma(R)$ is triangular, if R is one of the following.

- i) $Z_2 \oplus Z_2 \oplus Z_2$
- ii) $Z_4 \oplus Z_2$
- iii) $Z_4[x]/(x^2 + x + 1)$

Proof. A zero divisor graph is triangular if $\Gamma(R) = K_3$. Let R be a ring given by the Construction. Then $\Gamma(R) = K_{p^{hr}-1}$ or $\Gamma(R) = K_{p^{(h+1)r}-1}$. It suffices to find values of p, r and h for which $p^{hr} - 1$ or $p^{(h+1)r-1}$ equals 3. Now, let R be of characteristic $p = 2$. Then $\Gamma(R) = K_3$ when, $r = 1$ and $h = 2$. When R is of characteristic p^2 , then the graph is triangular when $p = 2, r = 1, h = 1$ and $p = 2, r = 2, h = 0$.

5 Rings of characteristic $p^k, k \geq 3$

Proposition 5.1 Let $char R = p^k, k \geq 3$.

$$\text{Then } \Gamma(R) = \begin{cases} p^{\binom{k+h}{2}r} - \text{partite} & \text{if } k \text{ is even} \\ p^{\binom{k-1+2h}{2}r} - \text{partite} & \text{if } k \text{ is odd} \end{cases}$$

Proof. Let $\lambda_1, \dots, \lambda_r \in R'$ with $\lambda_1 = 1$ such that $\overline{\lambda_1}, \dots, \overline{\lambda_r} \in R'/pR'$ form a basis for R'/pR' regarded as a vector space over its prime subfield F_p . Since the two cases do not overlap, we treat them in turn.

Case 1 : k is an even integer

Let $X_{i,s} = \{\sum_{i=1}^r a_i \lambda_i + \sum_{s=1}^h \lambda_s u_s\}, 1 \leq i \leq r, a_i \in \{0, j(\frac{k}{p^2})\}, 1 \leq j \leq \frac{k}{p^2} - 1, 1 \leq s \leq h$. Then $Z(R)^*$ is partitioned into the following mutually disjoint subsets, $V_{\sum a_i \lambda_i + \sum \lambda_s u_s} = X_{i,s} \setminus \{0\}, V_1 = Z(R)^* \setminus \bigcup_{i,s} V_{\sum a_i \lambda_i + \sum \lambda_s u_s}$.



These subsets are clearly nonempty and each contains nonadjacent vertices. Moreover, $Z(R)^* = V_1 \cup \bigcup_{i,s} \{V_{\sum a_i \lambda_i + \sum \lambda_s u_s}\}$. Now $|\bigcup_{i,s} \{V_{\sum a_i \lambda_i + \sum \lambda_s u_s}\}| = p^{\binom{k}{2}r} - 1$ and $|V_1| = 1$. So $|Z(R)^*| = p^{\binom{k}{2}r} - 1$ which implies that $\Gamma(R)$ is $p^{\binom{k}{2}r} - 1$ partite.

Case II: k is an odd integer

Let $X_{i,s} = \{\sum_{i=1}^r a_i \lambda_i + \sum_{s=1}^h \lambda_s u_s, 1 \leq i \leq r, a_i \in \{0, (j-1)p^{\frac{k+1}{2}}, 2 \leq j \leq p^{\frac{k-1}{2}}, 1 \leq s \leq h\}$. Then $Z(R)^{\hat{a}}$ is partitioned into the following mutually disjoint subsets, $V_{\sum a_i \lambda_i + \sum \lambda_s u_s} = X_{i,s} \setminus \{0\}, V_1 = Z(R)^{\hat{a}} \setminus \bigcup_{i,s} X_{i,s}$. The subsets are nonempty and each contains nonadjacent vertices. So $Z(R)^* = V_1 \cup \bigcup_{i,s} \{V_{\sum a_i \lambda_i + \sum \lambda_s u_s}\}$. Now $|\bigcup_{i,s} \{V_{\sum a_i \lambda_i + \sum \lambda_s u_s}\}| = p^{\binom{k-1}{2}r} - 1, |V_1| = 1$. So $|Z(R)^*| = |V_1| + |\bigcup_{i,s} V_{\sum a_i \lambda_i + \sum \lambda_s u_s}| = p^{\binom{k-1+2h}{2}r}$ showing that $\Gamma(R)$ is $p^{\binom{k-1+2h}{2}r}$ partite.

Corollary 5.2 Let $\text{char } R = p^k, k \geq 3$, then

i) $\text{diam}(\Gamma(R)) = 2$

ii) $\text{gr}(\Gamma(R)) = 3$

$$\text{iii) } b(\Gamma(R)) = \begin{cases} \frac{p^{\binom{k}{2}r} - 2}{p^{(k-1+h)r} - p^{\binom{k}{2}r}}, & \text{for } k \text{ is even} \\ \frac{p^{\binom{k-1}{2}r} - 1}{p^{(k-1+h)r} - p^{\binom{k-1}{2}r}}, & \text{for } k \text{ is odd} \end{cases}$$

Proof.

i) The annihilator, $\text{Ann}_{R'/pR'}(Z(R)) = (Z(R))^{n-1} \oplus U$. So the zero divisor of the form $p^{n-1}r' + \sum \lambda_i u_i$ is adjacent to every other nonzero zero divisor. Meanwhile, when $l+t \neq 0 \pmod{n}$, $r', s' \in R'$, then zero divisors of the form $p^l r' + \sum_{i=1}^h \lambda_i u_i$ and $p^t s' + \sum_{i=1}^h \lambda_i u_i$ are non adjacent. Thus $\text{diam}(\Gamma(R)) = 2$.

ii) The order of $\text{Ann}_{R'/pR'}(Z(R)) = p^{(h+1)r}$. So every zero divisor graph of R contains $K_{p^{(h+1)r}}$ subgraph. The invariants reveal that the least polygon in $\Gamma(R)$ is K_3

iii) Let k be an even integer. From the definition of $V_1, N(V_1) = V_{\sum a_i \lambda_i}$. So

$$|N(V_1)| = p^{\binom{k}{2}r} - 2. \text{ Also}$$

$$|V_1| = |Z(R)^{\hat{a}}| - |V_{\sum a_i \lambda_i + \sum \lambda_s u_s}| = p^{(k-1+h)r} - 1 - \left(p^{\binom{k}{2}r} - 2 \right) = p^{(k-1+h)r} - p^{\binom{k}{2}r} + 1.$$

If k is odd, then $|N(V_1)| = |V_{\sum a_i \lambda_i + \sum \lambda_s u_s}| = p^{\binom{k-1}{2}r} - 1$. Also observe that

$$|V_1| = |Z(R)^{\hat{a}} - X| = |Z(R)^{\hat{a}}| - |X| = p^{(k-1+h)r} - 1 - \left(p^{\binom{k-1}{2}r} - 1 \right) = p^{(k-1+h)r} - p^{\binom{k-1}{2}r}$$



. The ratio $|N(V_1)|/|V_1|$ completes the proof.

Proposition 5.3 There exists no ring in the Construction whose zero divisor graph, $\Gamma(R)$ is an n -gon, $n > 3$.

Proof. Suppose R is not a field and the cardinality of the vertices of nonzero zero divisors, $|V(\Gamma(R))| > 3$. Since $Z(R)$ is a nilpotent ideal, $Ann(Z(R)) \neq (0)$. Consider $0 \neq a \in Ann(Z(R))$, then a is adjacent to every other $b \in V(\Gamma(R))$. This completes the proof.

Proposition 5.4 Let R be a commutative ring given by the Construction. Then $diam(\Gamma(R)) = 0, 1$ or 2 .

Proof. If $R = \mathbb{Z}_2 \oplus \mathbb{Z}_2$, then $|Z(R)^*| = 1$; So $diam(\Gamma(R)) = 0$.

If $R = \mathbb{Z}_3 \oplus \mathbb{Z}_3$, then $|Z(R)^*| = 2$, so $diam(\Gamma(R)) = 1$.

For all the other constructed commutative rings, $diam(\Gamma(R)) = 2$ because $0 \neq a \in Ann(Z(R))$ is adjacent to every other vertex in $V(Z(R))$.

Proposition 5.5 The girth $gr(\Gamma(R)) = \infty$ if R is one of the following rings.

- i) $R = \mathbb{Z}_p[x] \langle f(x) \rangle$, where $f(x)$ is a monic irreducible polynomial modulo p of degree r .
- ii) $R = \mathbb{Z}_2 \oplus \mathbb{Z}_2$.
- iii) $R = \mathbb{Z}_3 \oplus \mathbb{Z}_3$.

Proof. In case (i), there exists no nonzero zero divisor.

In case (ii), there exists only one nonzero zero divisor, that is $(0,1)$.

For (iii), the ring has two adjacent nonzero zero divisors, that is $(0,1)$ and $(0,2)$. Cases (iv) and (v) are easy.

The next two results, summarize our findings on the zero divisor graphs of the equivalence classes of the rings considered in this paper.

Proposition 5.6 Let R be a ring given by the Construction. Then

$$\Gamma_E(R) = \begin{cases} \frac{k}{2} - \text{partite} & \text{if } k \text{ is even} \\ \frac{k+1}{2} - \text{partite} & \text{, if } k \text{ is odd} \end{cases}$$

Case I k is even.

$\Gamma_E(R)$ is partitioned into the following subsets

$$V_1 = \{(Z(R))^l \mid 1 \leq l \leq \frac{k}{2}\}$$

$V_j = \{(Z(R))^j\}, \frac{k}{2} < j \leq k-1$. For each $j, V_1 \cap V_j = \emptyset$ and V_j are mutually disjoint. Moreover

$V_1 \cup \bigcup_{j=\frac{k}{2}}^{k-1} \{V_j\} = \Gamma_E(R)$. The result follows by counting the disjoint subsets.

Case II: k is odd.



$\Gamma_E(R)$ is partitioned into the following

Proof. If $k = 1$ or 2 , then $\Gamma_E(R)$ is 1 -partite, so the result holds trivially.

For $k \geq 3$, we partition $\Gamma_E(R)$ into the following subsets

$$V_1 = \{(Z(R))^l \mid 1 \leq l \leq \frac{k-1}{2}\}$$

$$V_j = \{(Z(R))^j \mid \frac{k-1}{2} < j < k-1\}.$$

Then clearly for each j , $V_1 \cap V_j = \emptyset$ and V_j are mutually disjoint. Moreover $V_1 \cup \bigcup_{j=\frac{k-1}{2}}^{k-1} V_j = \Gamma_E(R)$. The result follows by counting the disjoint subsets.

Proposition 5.7 Let R be a ring given by the Construction in Section 2. Then,

- i) diameter, $diam(\Gamma_E(R)) = 2$.
- ii) girth, $gr(\Gamma_E(R)) = 3$.
- iii) binding number,

$$b(\Gamma_E(R)) = \begin{cases} \frac{k-4}{k} & \text{if } k \text{ is even} \\ \frac{k-5}{k-1} & \text{if } k \text{ is odd} \end{cases}$$

Proof. (i) and (ii) are easy to see.

To prove (iii), we consider the two separate cases when k is even and when k is odd respectively.

If k is even, $|V_1| = \frac{k}{2}$ while $|N(V_1)| = \frac{k-4}{2}$, so that $b(\Gamma_E(R)) = \frac{k-4}{k}$

Finally, when k is odd, $|V_1| = \frac{k-1}{2}$ while $|N(V_1)| = \frac{k-5}{2}$. Then $\frac{|N(V_1)|}{|V_1|} = \frac{k-5}{k-1}$

6. Conclusion

In this paper, we investigated the graphical properties of the zero divisors of the rings described by the Construction in Section 2. We recommend further research on the zero divisor graphs of the rings of idealizations and their automorphisms.

References

- [1] Abumoslem, A., Mohammad, E and Saeed, H. Some properties of zero divisor graphs, *International Mathematical Forum*, **7**(16), (2012), 761-767
- [2] Anderson, D.D., Naseer, M, Beck's coloring of a commutative ring, *Journal of Algebra*, **159**, (1993), 500-514.
- [3] Anderson, D.F., Livingston, P.S., The zero divisor graph of a commutative ring, *Journal of Algebra*, **217**(2), (1999), 434-447.
- [4] Diestel, R., "Graph Theory", Springer-Verlag, New York, 1997.
- [5] Sankeetha, S., The binding number of a zero divisor graph, *International Journal of Algebra*, **7**(5), (2013), 229-236.