



DSPACE

<https://dspace.org/>

**Intersection matrices associated with non trivial suborbit
corresponding to the action of rank 3 groups on the set
of unordered pairs**

Chepkorir, Betty; Rotich, John K; Tonui, Benard C; Langat, Reuben C

2014-12-12

Research gate

<http://ir-library.kabianga.ac.ke/handle/123456789/266>

BettyChepkorir, J. K. R., Tonui, B. C., & Langat, R. Intersection Matrices Associated With Non Trivial Suborbit Corresponding To The Action Of Rank 3 Groups On The Set Of Unordered Pairs.

Intersection Matrices Associated With Non Trivial Suborbit Corresponding To The Action Of Rank 3 Groups On The Set Of Unordered Pairs

BettyChepkorir, John K. Rotich, Benard C. Tonui, ReubenC. Langat

Abstract: In this paper we find intersection numbers and intersection matrices associated with each non trivial sub orbit corresponding to the action of rank 3 groups; The symmetric group S_5 , alternating group A_5 and The dihedral group D_5 on the set of unordered pairs. We showed that the column sum of the intersection matrices associated with Δ_i is equal the length of the suborbit Δ_i . They are also square matrices and of order 3×3 .

Index Terms: Intersection Matrices, Non Trivial Suborbit, Action of Rank 3 Groups, Set of Unordered Pairs

1 INTRODUCTION

In 1964, Higman introduced the rank of a group when he worked on finite permutation groups of rank 3. He showed that if G is a group acting transitively on a set X , where $|X| = n$ and if G is a rank 3 group of degree $n = k^2 + 1$, where k is the length of a G_x -orbit, $x \in X$ then $n = 5, 10, 50$ or 3250 . In 1970, Higman calculated the rank and subdegrees of the symmetric group S_n acting on a 2 element subsets from the set $X = \{1, 2, \dots, n\}$. He showed that the rank is 3 and the subdegrees are $1, 2(n-2), \binom{n-2}{2}$. In 1972, Cameron worked on suborbits of multiply permutation groups and later in 1974, he studied suborbits of primitive groups. In 1978, he dealt with the orbits of permutation groups of unordered pairs. In 1977, Neuman extended the work of Higman and Sims to finite permutation groups, edge coloured graphs and also matrices.

2 INTERSECTION MATRICES ASSOCIATED WITH THE ACTION OF $G = S_5$ ON $X^{(2)}$.

2.1 Intersection matrix corresponding to $\Delta_1 \{1, 2\}$.

Taking a $\Delta = \{1, 2\}$ in $X^{(2)}$ and $G_{\{1, 2\}}$ - orbits arranged as follows

$$\Delta_0 \{1, 2\} = \{1, 2\}$$

$$\Delta_1 \{1, 2\} = \{\{1, 3\}, \{1, 4\}, \{1, 5\}, \{2, 3\}, \{2, 4\}, \{2, 5\}\}$$

$$\Delta_2 \{1, 2\} = \{\{3, 4\}, \{3, 5\}, \{4, 5\}\}$$

We therefore arrange G_b - orbits as follows

$$\Delta_0 \{1, 3\} = \{1, 3\}$$

$$\Delta_1 \{1, 3\} = \{\{1, 2\}, \{1, 4\}, \{1, 5\}, \{3, 2\}, \{3, 4\}, \{3, 5\}\}$$

$$\Delta_2 \{1, 3\} = \{\{2, 4\}, \{2, 5\}, \{4, 5\}\}$$

$$\Delta_0 \{3, 4\} = \{3, 4\}$$

$$\Delta_1 \{3, 4\} = \{\{3, 1\}, \{3, 2\}, \{3, 5\}, \{4, 1\}, \{4, 2\}, \{4, 5\}\}$$

$$\Delta_2 \{3, 4\} = \{\{2, 1\}, \{2, 5\}, \{1, 5\}\}$$

From definition 6, the intersection numbers relative to the suborbit $\Delta_1 \{1, 2\}$ are defined by

$$\mu_{ij}^{(l)} = |\Delta_i(b) \cap \Delta_j \{1, 2\}|, \quad b \in \Delta_j \{1, 2\},$$

Hence we find the intersection numbers relative to $\Delta_1 \{1, 2\}$ as

- Betty Chepkoriris a lecturer at University of Kabianga, Kenya PH-+254728142051.
E-mail: bettyrono111@gmail.com
- John K. Rotich is a lecturer at University of Kabianga, Kenya and pursuing PhD degree program in Applied Mathematics in University of Eldoret, Kenya, PH-+254721273327.
E-mail: kimrott@gmail.com
- Benard C. Tonui, is a lecturer at University of Kabianga, Kenya
- Reuben C. Langatis a lecturer at University of Kabianga, Kenya

follows

$$\mu_{00}^{(1)} = |\Delta_1(\{1,2\}) \cap \Delta_0(\{1,2\})| = 0$$

$$\mu_{10}^{(1)} = |\Delta_1(\{1,2\}) \cap \Delta_1(\{1,2\})| = 6$$

$$\mu_{20}^{(1)} = |\Delta_1(\{1,2\}) \cap \Delta_2(\{1,2\})| = 0$$

$$\mu_{01}^{(1)} = |\Delta_1(\{1,3\}) \cap \Delta_0(\{1,2\})| = 1$$

$$\mu_{11}^{(1)} = |\Delta_1(\{1,3\}) \cap \Delta_1(\{1,2\})| = 3$$

$$\mu_{21}^{(1)} = |\Delta_1(\{1,3\}) \cap \Delta_2(\{1,2\})| = 2$$

$$\mu_{02}^{(1)} = |\Delta_1(\{3,4\}) \cap \Delta_0(\{1,2\})| = 0$$

$$\mu_{12}^{(1)} = |\Delta_1(\{3,4\}) \cap \Delta_1(\{1,2\})| = 4$$

$$\mu_{22}^{(1)} = |\Delta_1(\{3,4\}) \cap \Delta_2(\{1,2\})| = 2$$

By definition 7 the intersection matrix $M_1 = (\mu_{ij}^{(1)})_{i,j}$, associated with $\Delta_1\{1,2\}$ where $\mu_{ij}^{(1)}$ are the intersection numbers relative to $\Delta_1\{1,2\}$ is obtained as follows;

$$M_1 = \begin{bmatrix} \mu_{00}^{(1)} & \mu_{01}^{(1)} & \mu_{02}^{(1)} \\ \mu_{10}^{(1)} & \mu_{11}^{(1)} & \mu_{12}^{(1)} \\ \mu_{20}^{(1)} & \mu_{21}^{(1)} & \mu_{22}^{(1)} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 1 & 0 \\ 6 & 3 & 4 \\ 0 & 2 & 2 \end{bmatrix}$$

2.2 Intersection matrix corresponding to $\Delta_2\{1,2\}$

From definition 6, the intersection numbers relative to the suborbit $\Delta_2\{1,2\}$ are defined by

$$\mu_{ij}^{(2)} = |\Delta_2(b) \cap \Delta_i\{1,2\}|, \quad b \in \Delta_j\{1,2\},$$

We therefore find the intersection numbers relative to

$\Delta_2\{1,2\}$ as follows

$$\mu_{00}^{(2)} = |\Delta_2(\{1,2\}) \cap \Delta_0(\{1,2\})| = 0$$

$$\mu_{10}^{(2)} = |\Delta_2(\{1,2\}) \cap \Delta_1(\{1,2\})| = 0$$

$$\mu_{20}^{(2)} = |\Delta_2(\{1,2\}) \cap \Delta_2(\{1,2\})| = 3$$

$$\mu_{01}^{(2)} = |\Delta_2(\{1,3\}) \cap \Delta_0(\{1,2\})| = 0$$

$$\mu_{11}^{(2)} = |\Delta_2(\{1,3\}) \cap \Delta_1(\{1,2\})| = 2$$

$$\mu_{21}^{(2)} = |\Delta_2(\{1,3\}) \cap \Delta_2(\{1,2\})| = 1$$

$$\mu_{02}^{(2)} = |\Delta_2(\{3,4\}) \cap \Delta_0(\{1,2\})| = 1$$

$$\mu_{12}^{(2)} = |\Delta_2(\{3,4\}) \cap \Delta_1(\{1,2\})| = 2$$

$$\mu_{22}^{(2)} = |\Delta_2(\{3,4\}) \cap \Delta_2(\{1,2\})| = 0$$

By definition 7 the intersection matrix $M_2 = (\mu_{ij}^{(2)})_{i,j}$,

associated with $\Delta_2\{1,2\}$ where $\mu_{ij}^{(2)}$ are the intersection numbers relative to $\Delta_2\{1,2\}$ is obtained as follows;

$$M_2 = \begin{bmatrix} \mu_{00}^{(2)} & \mu_{01}^{(2)} & \mu_{02}^{(2)} \\ \mu_{10}^{(2)} & \mu_{11}^{(2)} & \mu_{12}^{(2)} \\ \mu_{20}^{(2)} & \mu_{21}^{(2)} & \mu_{22}^{(2)} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 1 \\ 0 & 2 & 2 \\ 3 & 1 & 0 \end{bmatrix}$$

2.3 Properties of the intersection matrices associated with $\Delta_1\{1,2\}$ and $\Delta_2\{1,2\}$

The column sum of the intersection matrix associated with Δ_i

is equal to the length of the suborbit Δ_i . We can see that the

column sum of M_1 is 6 also the column sum of M_2 is 3. M_1 and

M_2 are square matrices of order 3

3 INTERSECTION MATRICES ASSOCIATED WITH THE ACTION OF $G = A_5$ ON $X^{(2)}$

3.1 Intersection matrix corresponding to $\Delta_1 \{1, 2\}$

we take $a = \{1, 2\}$ in $X^{(2)}$ and $G_{\{1, 2\}}$ - orbits arranged as follows

$$\Delta_0 \{1, 2\} = \{1, 2\}$$

$$\Delta_1 \{1, 2\} = \{\{1, 3\}, \{1, 4\}, \{1, 5\}, \{2, 3\}, \{2, 4\}, \{2, 5\}\}$$

$$\Delta_2 \{1, 2\} = \{\{3, 4\}, \{3, 5\}, \{4, 5\}\}$$

We therefore arrange G_b - orbits as follows

$$\Delta_0 \{1, 3\} = \{1, 3\}$$

$$\Delta_1 \{1, 3\} = \{\{1, 2\}, \{1, 4\}, \{1, 5\}, \{3, 2\}, \{3, 4\}, \{3, 5\}\}$$

$$\Delta_2 \{1, 3\} = \{\{2, 4\}, \{2, 5\}, \{4, 5\}\}$$

$$\Delta_0 \{3, 4\} = \{3, 4\}$$

$$\Delta_1 \{3, 4\} = \{\{3, 1\}, \{3, 2\}, \{3, 5\}, \{4, 1\}, \{4, 2\}, \{4, 5\}\}$$

$$\Delta_2 \{3, 4\} = \{\{2, 1\}, \{2, 5\}, \{1, 5\}\}$$

From definition 1.1.6.1, the intersection numbers relative to the

suborbit $\Delta_1 \{1, 2\}$ are defined by

$$\mu_{ij}^{(1)} = |\Delta_i(b) \cap \Delta_j \{1, 2\}|, \quad b \in \Delta_j \{1, 2\},$$

Hence we find the intersection numbers relative to $\Delta_1 \{1, 2\}$ as

follows

$$\mu_{00}^{(1)} = |\Delta_1(\{1, 2\}) \cap \Delta_0(\{1, 2\})| = 0$$

$$\mu_{10}^{(1)} = |\Delta_1(\{1, 2\}) \cap \Delta_1(\{1, 2\})| = 6$$

$$\mu_{20}^{(1)} = |\Delta_1(\{1, 2\}) \cap \Delta_2(\{1, 2\})| = 0$$

$$\mu_{01}^{(1)} = |\Delta_1(\{1, 3\}) \cap \Delta_0(\{1, 2\})| = 1$$

$$\mu_{11}^{(1)} = |\Delta_1(\{1, 3\}) \cap \Delta_1(\{1, 2\})| = 3$$

$$\mu_{21}^{(1)} = |\Delta_1(\{1, 3\}) \cap \Delta_2(\{1, 2\})| = 2$$

$$\mu_{02}^{(1)} = |\Delta_1(\{3, 4\}) \cap \Delta_0(\{1, 2\})| = 0$$

$$\mu_{12}^{(1)} = |\Delta_1(\{3, 4\}) \cap \Delta_1(\{1, 2\})| = 4$$

$$\mu_{22}^{(1)} = |\Delta_1(\{3, 4\}) \cap \Delta_2(\{1, 2\})| = 2$$

By definition 1.1.6.2 the intersection matrix $M_1 = (\mu_{ij}^{(1)})_{i,j}$,

associated with $\Delta_1 \{1, 2\}$ where $\mu_{ij}^{(1)}$ are the intersection

numbers relative to $\Delta_1 \{1, 2\}$ is obtained as follows;

$$M_1 = \begin{bmatrix} \mu_{00}^{(1)} & \mu_{01}^{(1)} & \mu_{02}^{(1)} \\ \mu_{10}^{(1)} & \mu_{11}^{(1)} & \mu_{12}^{(1)} \\ \mu_{20}^{(1)} & \mu_{21}^{(1)} & \mu_{22}^{(1)} \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 6 & 3 & 4 \\ 0 & 2 & 2 \end{bmatrix}$$

3.2 Intersection matrix corresponding to $\Delta_2 \{1, 2\}$

From definition 1.1.6.1, the intersection numbers relative to the

suborbit $\Delta_2 \{1, 2\}$ are defined by

$$\mu_{ij}^{(2)} = |\Delta_2(b) \cap \Delta_j \{1, 2\}|, \quad b \in \Delta_j \{1, 2\},$$

We therefore find the intersection numbers relative to

$\Delta_2 \{1, 2\}$ as follows

$$\mu_{00}^{(2)} = |\Delta_2(\{1, 2\}) \cap \Delta_0(\{1, 2\})| = 0$$

$$\mu_{10}^{(2)} = |\Delta_2(\{1, 2\}) \cap \Delta_1(\{1, 2\})| = 0$$

$$\mu_{20}^{(2)} = |\Delta_2(\{1, 2\}) \cap \Delta_2(\{1, 2\})| = 3$$

$$\mu_{01}^{(2)} = |\Delta_2(\{1, 3\}) \cap \Delta_0(\{1, 2\})| = 0$$

$$\mu_{11}^{(2)} = |\Delta_2(\{1, 3\}) \cap \Delta_1(\{1, 2\})| = 2$$

$$\mu_{21}^{(2)} = |\Delta_2(\{1, 3\}) \cap \Delta_2(\{1, 2\})| = 1$$

$$\mu_{02}^{(2)} = |\Delta_2(\{3, 4\}) \cap \Delta_0(\{1, 2\})| = 1$$

$$\mu_{12}^{(2)} = |\Delta_2(\{3, 4\}) \cap \Delta_1(\{1, 2\})| = 2$$

$$\mu_{22}^{(2)} = |\Delta_2(\{3, 4\}) \cap \Delta_2(\{1, 2\})| = 0$$

By definition 1.1.6.2 the intersection matrix $M_2 = (\mu_{ij}^{(2)})_{i,j}$,

associated with $\Delta_2\{1, 2\}$ where $\mu_{ij}^{(2)}$ are the intersection

numbers relative to $\Delta_2\{1, 2\}$ is obtained as follows;

$$M_2 = \begin{bmatrix} \mu_{00}^{(2)} & \mu_{01}^{(2)} & \mu_{02}^{(2)} \\ \mu_{10}^{(2)} & \mu_{11}^{(2)} & \mu_{12}^{(2)} \\ \mu_{20}^{(2)} & \mu_{21}^{(2)} & \mu_{22}^{(2)} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 2 & 2 \\ 3 & 1 & 0 \end{bmatrix}$$

3.3 Properties of the intersection matrices associated with $\Delta_1\{1, 2\}$, and $\Delta_2\{1, 2\}$.

The column sum of the intersection matrix associated with Δ_i

is equal to the length of the suborbit Δ_i . We can see that the

column sum of M_1 is 6 also the column sum of M_2 is 3. M_1 and

M_2 are square matrices of order 3

4 INTERSECTION MATRICES ASSOCIATED WITH THE ACTION OF $G=D_5$ ON X

By Definition 6, given an arrangement of the G_a -orbits, the

G_b -orbits are arranged such that if $b \in X$ and $g(a) = b$ then,

$$g(\Delta_i(a)) = \Delta_i(g(b)) = \Delta_i(b)$$

4.1 Intersection matrix corresponding to $\Delta_1(1)$.

Taking $a = 1$ in X and G_1 -orbits arranged as follows,

$$\Delta_0(1) = \{1\}.$$

$$\Delta_1(1) = \{2, 5\},$$

$$\Delta_2(1) = \{3, 4\},$$

We arrange the G_b -orbits as follows:

$$\Delta_0(2) = \{2\}.$$

$$\Delta_1(2) = \{1, 3\},$$

$$\Delta_2(2) = \{4, 5\},$$

$$\Delta_0(3) = \{3\}.$$

$$\Delta_1(3) = \{1, 5\},$$

$$\Delta_2(3) = \{2, 4\},$$

From definition 6, the intersection numbers relative to the suborbit $\Delta_1(1)$ are defined by

$$\mu_{ij}^{(1)} = |\Delta_i(b) \cap \Delta_j(1)|, \quad b \in \Delta_j(1),$$

Hence we find the intersection numbers relative to $\Delta_1(1)$ as follows

$$\mu_{00}^{(1)} = |\Delta_1(1) \cap \Delta_0(1)| = 0$$

$$\mu_{10}^{(1)} = |\Delta_1(1) \cap \Delta_1(1)| = 2$$

$$\mu_{20}^{(1)} = |\Delta_1(1) \cap \Delta_2(1)| = 0$$

$$\mu_{01}^{(1)} = |\Delta_1(2) \cap \Delta_0(1)| = 1$$

$$\mu_{11}^{(1)} = |\Delta_1(2) \cap \Delta_1(1)| = 0$$

$$\mu_{21}^{(1)} = |\Delta_1(2) \cap \Delta_2(1)| = 1$$

$$\mu_{02}^{(1)} = |\Delta_1(3) \cap \Delta_0(1)| = 1$$

$$\mu_{12}^{(1)} = |\Delta_1(3) \cap \Delta_1(1)| = 1$$

$$\mu_{22}^{(1)} = |\Delta_1(3) \cap \Delta_2(1)| = 0$$

By definition 7 the intersection matrix $M_1 = (\mu_{ij}^{(1)})_{i,j}$,

associated with $\Delta_1 \{1, 2\}$ where $\mu_{ij}^{(1)}$ are the intersection numbers relative to $\Delta_1(1)$ is obtained as follows;

$$M_1 = \begin{bmatrix} \mu_{00}^{(1)} & \mu_{01}^{(1)} & \mu_{02}^{(1)} \\ \mu_{10}^{(1)} & \mu_{11}^{(1)} & \mu_{12}^{(1)} \\ \mu_{20}^{(1)} & \mu_{21}^{(1)} & \mu_{22}^{(1)} \end{bmatrix} = \begin{bmatrix} 0 & 1 & 1 \\ 2 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

4.2 Intersection matrix corresponding to $\Delta_2(1)$

From definition 1.1.6.1, the intersection numbers relative to the suborbit $\Delta_2(1)$ are defined by

$$\mu_{ij}^{(2)} = |\Delta_2(b) \cap \Delta_i(1)|, \quad b \in \Delta_j(1),$$

We therefore find the intersection numbers relative to $\Delta_2(1)$

as follows

$$\mu_{00}^{(2)} = |\Delta_2(1) \cap \Delta_0(1)| = 0$$

$$\mu_{10}^{(2)} = |\Delta_2(1) \cap \Delta_1(1)| = 0$$

$$\mu_{20}^{(2)} = |\Delta_2(1) \cap \Delta_2(1)| = 2$$

$$\mu_{01}^{(2)} = |\Delta_2(2) \cap \Delta_0(1)| = 0$$

$$\mu_{11}^{(2)} = |\Delta_2(2) \cap \Delta_1(1)| = 1$$

$$\mu_{21}^{(2)} = |\Delta_2(2) \cap \Delta_2(1)| = 1$$

$$\mu_{02}^{(2)} = |\Delta_2(3) \cap \Delta_0(1)| = 0$$

$$\mu_{12}^{(2)} = |\Delta_2(3) \cap \Delta_1(1)| = 1$$

$$\mu_{22}^{(2)} = |\Delta_2(3) \cap \Delta_2(1)| = 1$$

By definition 6 the intersection matrix $M_2 = (\mu_{ij}^{(2)})_{i,j}$,

associated with $\Delta_2(1)$ where $\mu_{ij}^{(2)}$ are the intersection

numbers relative to $\Delta_2(1)$ is obtained as follows;

$$M_2 = \begin{bmatrix} \mu_{00}^{(2)} & \mu_{01}^{(2)} & \mu_{02}^{(2)} \\ \mu_{10}^{(2)} & \mu_{11}^{(2)} & \mu_{12}^{(2)} \\ \mu_{20}^{(2)} & \mu_{21}^{(2)} & \mu_{22}^{(2)} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 2 & 1 & 1 \end{bmatrix}$$

4.3 Properties of the intersection matrices associated with $\Delta_2(1)$, and $\Delta_2(1)$.

The column sum of the intersection matrix associated with Δ_i is equal to the length of the suborbit Δ_i . We can see that the column sum of M_1 is 6 also the column sum of M_2 is 3. M_1 and M_2 are square matrices of order 3

5 CONCLUSION

We conclude that Intersection matrices associated with the action of rank 3 on $X^{(2)}$ are square matrices of order 3×3 and that the column sum of the intersection matrices associated with Δ_i is equal to the length of the suborbit Δ_i .

REFERENCES

- [1] Akbas, M. 2001. Suborbital graphs for modular group, Bulletin of the London mathematical society 33:647-652.
- [2] Burnside, W. 1911. Theory of groups of finite order, Cambridge University Press, Cambridge (Dover reprint 1955).
- [3] Bon, J. V. and Cohen, A.M. 1989. Linear groups and distance-transitive groups, European Journal of combinatorics 10:399-411.
- [4] Cameron, P. J. 1972. Permutation groups with multiply transitive suborbits I. Proc. London Math. Soc. 23 (3): 427 – 440.
- [5] Cameron, P. J. 1974. Permutation groups with multiply transitive suborbits II. Bull. London Math. Soc. 6: 136 - 140.
- [6] Cameron, P. J. 1978. Orbit of permutation groups on Unordered sets. J. London. Math. Soc. 17 (2): 410- 414.
- [7] Chartrand, G. 1993. Applied and algorithmic graph theory. International series in pure and applied Mathematics: 238-240.
- [8] Coxeter, H. S. M. 1986. My graph, proceedings of London mathematical society 46:117-136.
- [9] Faradžev, I. A. and Ivanov, A. A. 1990. Distance-transitive representations of group G with $PSL(2q) < G < P\Gamma L(2, q)$, European journal of combinatorics 11:347-356.
- [10] Harary, F. 1969. Graph Theory. Addison – Wesley Publishing Company, New York.
- [11] Higman, D. G. 1964. Finite permutation groups of rank 3. Math Zeitschrift 86: 145 – 156.
- [12] Higman, D. G. 1970. Characterization of families of rank 3 permutation groups by subdegrees. I. Arch. Math 21: 151-156.

- [13] Jones, G. A.; Singerman, D. and Wicks, K., 1991. Generalized Farey graphs in groups, St. Andrews 1989, Eds. C. Campbell and E.F. Robertson, London mathematical society lecture notes series 160, Cambridge University, Cambridge 316 – 338.
- [14] Kamuti, I. N. 1992. Combinatorial formulas, invariants and structures associated with primitive permutation representations of PSL (2, q) and PGL (2, q). Ph. D. Thesis, Southampton University, U.K.
- [15] Kamuti, I. N. 2006. Subdegrees of primitive permutation representation of PGL (2,q), East African journal of physical sciences 7(1/2):25-41.
- [16] Kangogo, M. 2008. Some properties of symmetric group S_6 and associated combinatorial formulas and structures, M.Sc. Project; Kenyatta, University, Kenya.
- [17] Krishnamurthy, V. 1985. Combinatorics, theory and applications, Affiliated East West Press Private Limited, New Delhi.
- [18] Neumann, P. M. 1977. Finite permutation groups edge-coloured graphs and matrices edited by M. P. J. Curran, Academic Press, London.
- [19] Petersen, J. 1898. Sur le Theore'me de Tait Interned Math 5:225-227.
- [20] Rose, J. S. 1978. A Course on group theory. Cambridge University Press, Cambridge.
- [21] Rosen, K. H. 1999. Handbook of Discrete and combinatorial Mathematics. CRC Press, New Jersey.
- [22] Rotman, J. J. 1973. The theory of groups: An introduction. Allyn and Bacon, Inc. Boston, U.S.A.
- [23] Rotich, K. S. 2008. Some properties of the symmetric group S_7 acting on ordered and unordered Pairs and the associated combinatorial structures, M.Sc. Project, Kenyatta University, Kenya.
- [24] Galois, C. 1830. Self-Conjugate sub-groups; simple and composite groups. Cambridge University Press Cambridge.
- [25] Jordan, C. 1870. Traite des substitutions et des Equations Algebriques. Gauthier – Villas, Paris.
- [26] Sims, C. C. 1967. Graphs and finite permutation groups. Math. Zeitschrift 95: 76 – 86.
- [27] Wielandt, H. 1964. Finite Permutation Groups. Academic Press, New York and London.

APPENDIX:

1. NOTATIONS

- i). S_n , Symmetric group of degree n and order $n!$.
- ii). $|G|$, The order of a group G .

- iii). $X^{(2)}$, the set of unordered pairs from the set $X \{1, 2, \dots, n\}$
- iv). $\{t, q\}$, Unordered pair.
- v). Δ_i , the i^{th} orbit or suborbit.
- vi). $\mu^{(i)}$, the intersection number relative to a suborbit $\Delta_i(a)$.
- vii). M_i , intersection matrix of a suborbit $\Delta_i(a)$.

2. DEFINITION AND PRELIMINARY RESULTS

Definition 1

The alternating group A_n is the subgroup of S_n comprising of all even permutations. Its order is $\frac{n!}{2}$.

Definition 2

Let X be a finite set $\{1, 2, \dots, n\}$, then a symmetric group of regular n -gon is called a dihedral group denoted by Δ_n .

Definition 3

When G act on a set X , X is divided into disjointed equivalence classes of the action called orbits. The orbits containing x is called the orbit of x , denoted by $\text{Orb}_G(x)$.

Definition 4

Let G be transitive on X and let G_x be the stabilizer of a point $x \in X$. The orbits $\Delta_0, \Delta_1, \Delta_2, \dots, \Delta_{r-1}$ of G_x on X are the suborbits of G .

Definition 5

The rank r of G is the number of the suborbits of G while the lengths of the suborbits of G are known as the subdegrees of G .

Note: The cardinalities of the suborbits Δ_i and are independent of the choice of $x \in X$.

INTERSECTION NUMBERS AND INTERSECTION MATRICES

Definition 6

Let G be a finite group acting on a finite set X and $\Delta_i(a)$ be the i^{th} G_a -orbit for $a \in X$ and for a given arrangement of the G_a -orbits. The G_b -orbit, $b \in X$, are also arranged such that $g(a) = b$, then $g(\Delta_i(a)) = \Delta_i(g(a)) = \Delta_i(b)$. The intersection numbers relative to a suborbit $\Delta_i(a)$ are defined by

$$\mu_{ij}^{(i)} = |\Delta_i(b) \cap \Delta_i(a)|, \quad (b \in \Delta_j(a))$$

Definition 7

If the rank of G is r , then the $r \times r$ matrix $M_i = (\mu_{ij}^{(i)})_{i,j}$ is

called the intersection matrix of $\Delta_l(a)$.

Theorem 1 [Higman, [11]]

If $|\Delta_i| = n_i$ and $\Delta_i^* = \Delta_{i^*}$ is the suborbit paired with Δ_i , then

$$\text{a) } \mu_{i0}^{(l)} = \begin{cases} n_i & \text{if } i = l \\ 0 & \text{if } i \neq l \end{cases}$$

$$\text{b) } \mu_{i0}^{(l)} = \begin{cases} 1 & \text{if } j = l^* \\ 0 & \text{if } j \neq l^* \end{cases}$$

$$\text{c) } n_j \mu_{ij}^{(l)} = n_i \mu_{ji}^{(l^*)} \text{ and } n_i \mu_{i^*i}^{(j)} = n_j \mu_{i^*j}^{(l)} = n_l \mu_{j^*l}^{(i)}$$