



DSPACE

<https://dspace.org/>

Promoting Agriculture through Data Analytics: Pathways to Strengthen Food Security.

Reuben Cheruiyot Lang'at

2026

American Journal of Applied Statistics and Economics (AJASE)


<https://ir-library.kabianga.ac.ke/handle/123456789/1150>



American Journal of Applied Statistics and Economics (AJASE)

ISSN: 2992-927X (ONLINE)

VOLUME 5 ISSUE 1 (2026)



PUBLISHED BY
E-PALLI PUBLISHERS, DELAWARE, USA

Promoting Agriculture through Data Analytics: Pathways to Strengthen Food Security.

Reuben Cheruiyot Lang^{at}*

Article Information

Received: November 09, 2025

Accepted: February 07, 2025

Published: March 19, 2026

Keywords

*Analysis Of Variance, Data
Analytic, Digital Agriculture.
Food Security, Precision
Agriculture, Regression Analysis,
Remote Sensing*

ABSTRACT

Globally, agriculture faces escalating pressures from climate change, population growth, declining soil fertility, and market volatility. As food insecurity intensifies, especially in Sub-Saharan Africa, data analytics has emerged as a transformative tool for improving agricultural decision-making, productivity, and resilience. This paper examines the role of data analytics in enhancing crop forecasting, optimizing resource use, improving extension services, and designing evidence-based policies to ensure sustainable food systems. Drawing from empirical studies and international development reports, the paper argues that data-driven agriculture provides an effective pathway for addressing chronic food insecurity while supporting national development strategies. This study presents four illustrative, reproducible analyses using built-in R datasets that highlight common analytical approaches such as descriptive statistics, regression analysis, and analysis of variance, and how their outputs can inform agricultural decisions. The paper concludes with policy recommendations for integrating analytics capacity into agricultural institutions, particularly in low and middle-income countries striving to modernize their food systems.

INTRODUCTION

World over, food security remains a pressing policy and development challenge. The Food and Agriculture Organization (FAO) estimates that tens to hundreds of millions of people remain food-insecure, and multiple shocks, including the recent COVID-19 pandemic, armed conflicts, trade disruptions, and climate extremes, continue to exacerbate the problem. Closing production gaps, crop and variety diversification, improving distribution, and building resilient value chains are essential, and the role of data analytics is increasingly taking its stage as a central toolset to accomplish these goals (FAO, SOFI).

Data analytics in agriculture which is often called digital agriculture, precision agriculture, or smart farming, covers a wide spectrum: remote sensing and satellite monitoring, in-field sensors and Internet of Things (IoT) devices, farm management systems, supply-chain traceability platforms, and advanced analytics (machine learning, optimization) applied to those data streams (Wolfert, Ge, Verdouw, & Bogaardt, 2017)(Wolfert *et al*, 2017). These technologies promise improved decision-making, resource efficiency, early warning for pests and diseases, and better targeting of interventions for vulnerable populations. At the same time, there are socio-economic, governance, and infrastructural barriers that can limit equitable realization of benefits.

This paper pursues three aims: (1) synthesize the academic and policy evidence on how data analytics can promote agricultural productivity and food security; (2) demonstrate, with concrete R examples using built-in datasets, how standard analytic techniques yield actionable insights for agriculture; and (3) provide practical recommendations for policy makers, development agencies, and researchers to accelerate inclusive adoption.

Conceptual framework: How analytics influences food security

Food security is commonly understood across four dimensions: availability, access, utilization, and stability. Data analytics can affect each dimension through specific mechanisms:- it is obvious that food has to be available to humanity. For this to be realized every nation has to know or at least be in a position to predict production and supply within a given season or period. For this to happen yield prediction, field-level recommendations, improved irrigation scheduling, and optimized input use, forms the basis for increased and effective supply. Satellite and ground sensors provide spatially explicit data to manage fields at sub-field resolution. (Wolfert *et al*, 2017)

Farmers on the other hand have to gain access to better market information systems and supply-chain analytics reduce post-harvest losses, improve price discovery, and enhance logistics, potentially lowering consumer prices and increasing market access for producers. Equally consumers need to have such information to plan and stock their foods at affordable prices. Significant improvement has been noted in food safety as a result of use of data analytics by enabling predictive modeling of contamination risks and enhancing supply chain traceability, while in nutrition, it has enabled personalized diet recommendations and public health monitoring. Traceability and quality monitoring as informed by data analytics enhances safer food handling and targeted nutrition interventions (Dogho & Babatunde, 2025).

Climate change and abrupt change of weather patterns can cause untold losses in agriculture. Various parts of the world have had their agricultural fields being covered or swept away during landslides and other sudden calamities such as earthquakes. It is also never a better option

¹ University of Kabianga P.O. Box 2030-20200 Kericho, Kenya

* Corresponding author's e-mail: rlangat@kabianga.ac.ke

when the rain subsides, the extreme case of prolonged droughts threaten food stability especially where no proper planning is in place. To have food stability and resilience to these shocks, early warning systems driven by climate and crop models, Geographical Information Systems (GIS) remote sensing, and risk analytics come in handy in not only stabilizing production and incomes but also providing the much needed information to enable proper proactive decision making (Yu, *et al.*, 2025; Wang, Akber, & Aziz, 2025).

This framework shows analytic interventions acting across scales right from smallholder plots to national monitoring systems and across actors such as farmers, cooperatives, extension services, agribusinesses, and policy makers among others.

LITERATURE REVIEW

Big data and smart farming

Wolfert *et al.* (2017) provide a widely cited review on how big data and interoperability are reshaping farm management systems and decision workflows. Their review emphasizes data heterogeneity, the value of integrating sensor and enterprise datasets, and the governance challenges of data ownership and privacy. With current trends in population pressure and climatic conditions that seem to undermine food production, it is inevitable that proper planning and accurate data analysis is necessary if production is to be stepped up to meet the ever increasing demand. Big data analytics plays a pivotal role in this transformation, enabling farmers to use real-time and historical data to predict outcomes, allocate resources effectively, and reduce risks (Lohit & Mujahid, 2022; Aissi, Benjelloun, Lakhri, & Ali, 2023).

Precision and digital agriculture: technologies and evidence

Recent reviews and empirical studies document the advances in precision agriculture that include the Global Positioning System (GPS)-guided application, variable rate fertilization, Unmanned Aerial Vehicle (UAV) or satellite crop monitoring and the emerging evidence of yield increases and input savings. However, outcomes are context dependent and depend heavily on extension, data literacy, and access to affordable technology. These technological advances are playing a central role in the transformation of agricultural practices, offering solutions to optimize inputs, increase productivity, and mitigate environmental impact (Soussi, Zero, Sacile, Trincherro, & Fossa, 2025). In another dimension emergence more sophisticated technologies such as robotics, artificial intelligence (AI), sensors, and Internet of Things (IoT) devices have been used to perform farming tasks with minimal human help. This has led to what is termed as Agricultural automation (AA). The automation aims to enhance the sustainability, efficiency, and productivity of farming operations. AA is a result of the revolutionary way that AI has emerged to solve these issues and improve agricultural processes. AI-driven apps assess the health

of the soil by analyzing data from soil samples and give information on factors such as soil pH, moisture content, nutrient content, and organic matter among others. This way, if farmers incorporate the use of AA, a precision-based, data-driven method, it would be expected that farming would become more sustainable, productive, and profitable as opposed to the use of traditional approaches to agriculture (Sarker, Abdolrasol, Md, Kadir, Ahmad, & Olazagoitia, 2025).

Remote sensing and large scale monitoring

The technology of remote sensing enables data collection, procession, and analysis of remotely sensed data. It also makes it possible to retrieve, synthesize, and visualize valuable geospatial information for use in agriculture. In particular, remote sensing technology empowers capability for large scale field level or regional assessment and monitoring of crop land cover, crops growth and condition via remotely sensed data from satellite or aerial sensors.

The product such as Normalized Difference Vegetation Index (NDVI), a measure used to detect and monitor vegetation health and density using satellite or aerial imagery. It compares the reflectance of near-infrared light to red light. Healthy vegetation is known to reflect strongly on infrared light while it absorbs that of red light. The resulting index ranges from -1 to +1, with values closer to 1 indicating dense, healthy vegetation, and values near 0 or negative indicating non-vegetated surfaces like water, snow, or barren ground. To determine the wetness or dryness of a place, soil moisture proxies are analyzed. These are measurements that estimate soil moisture levels, especially over large areas where direct measurement is not feasible. Common proxies include the Topographic Witness Index (TWI) and drought indices like the Z-index and Component Moisture Index (CMI). TWI uses topography to predict where water will accumulate, while drought indices are derived from precipitation and other variables. These indices are critical for national-level monitoring, drought early warning, and yield forecasting. Reviews show steady improvements in data availability (e.g., Sentinel, Landsat) and algorithms for crop classification and area estimation, but also highlight challenges in ground truthing and resolution for fragmented smallholder landscapes (Yang, Wu, Di, & Üstündağ, 2017) (Yang, Wu, Di, & Üstündağ, 2017)

Policy landscape and digital agriculture strategies

International institutions (World Bank, FAO) emphasize the need for national digital agriculture roadmaps, interoperability standards, and investment in connectivity and data governance. The World Bank's Digital Agriculture Roadmap Playbook offers practical guidance for governments designing digital transformation strategies in agriculture. A Digital Agriculture Roadmap (DAR) is a strategy, investment, and implementation plan for a country or region, intended to guide transformation of the agriculture sector using digital technologies. It outlines key role played by policymakers

and funders in enabling countries to realize the benefits of the DAR approach. These include developing and institutionalizing DARs within national and regional agriculture, digital, mobilizing funds for implementation, by allocating domestic budgets and donor funds where possible, coordinating across the ecosystem to maximize resource efficiency and effectiveness and ensuring strong ownership and institutional capacity to lead planning and implementation, as well as investing in digital skills development (World Bank, 2025).

While the literature shows strong technical promise, two recurring barriers are: (1) unequal access and capacity (digital divide); and (2) institutional and governance gaps (data sharing, standards, financing). Addressing these is essential for analytics to create equitable food-security gains.

MATERIALS AND METHODS

Research design and analytical approach

This study adopts a mixed analytical design, combining a systematic narrative review of literature with illustrative quantitative analyses to demonstrate how data analytics can inform agricultural decision-making and food security outcomes. The methodological approach is explanatory and demonstrative rather than inferential for population estimation. The objective is not to generalize from the illustrative datasets to real-world agricultural systems, but rather to show how standard analytical techniques commonly used in agricultural research can generate insights that are directly relevant to productivity, resilience, and food security planning.

Data sources and justification

The empirical illustrations in this paper rely on built-in datasets available in base R, specifically the *swiss*, *Orange*, *CO₂*, and *PlantGrowth* datasets. These datasets were selected deliberately for pedagogical and methodological clarity. They are well-documented, widely used in statistical education and research, and allow for transparent replication of results. Although these datasets are not modern agricultural production datasets, they embody core structures and relationships that are directly analogous to real agricultural and food system data.

The *swiss* dataset represents socio-economic and demographic conditions across French-speaking Swiss provinces in the late nineteenth century. It is used to illustrate how regression analysis can identify socio-economic correlates of agricultural labor dependence, which is relevant for understanding structural transformation, rural livelihoods, and long-term food availability.

The *Orange* dataset contains longitudinal measurements of tree circumference for multiple orange trees observed over time. This dataset is used to demonstrate growth-curve analysis, which is conceptually analogous to crop growth monitoring, yield development, and phenological analysis in agriculture.

The *CO₂* dataset provides experimental data on carbon

dioxide uptake by plants under different concentrations, treatments, and origins. This dataset is used to illustrate how physiological response data can be analyzed to understand crop performance under environmental stress, a critical concern for climate-resilient agriculture and food stability.

The *PlantGrowth* dataset contains experimental measurements of plant biomass under different treatment conditions. It is employed to demonstrate experimental design analysis using one-way analysis of variance (ANOVA) and post-hoc comparisons, which are fundamental tools in agronomic trials evaluating fertilizers, soil amendments, or crop management practices.

Analytical procedures

All statistical analyses were conducted using R statistical software (R Core Team, 2024). The analysis proceeded in a stepwise manner aligned with common agricultural data analytics workflows.

Descriptive statistics were first computed to summarize central tendencies and variability within each dataset. Graphical methods, including line plots, scatterplots, and boxplots, were used extensively to visualize patterns, trends, and group differences. Visualization plays a central role in agricultural analytics by enabling intuitive interpretation of complex data by researchers, extension officers, and policy makers.

For multivariate relationship analysis, multiple linear regression was applied to the *swiss* dataset. Agriculture, measured as the percentage of males engaged in agricultural labor, was modeled as a function of education, examination performance, and infant mortality. Ordinary Least Squares (OLS) estimation was used, and model assumptions were evaluated through standard diagnostic plots, including residuals versus fitted values, normal Q-Q plots, scale-location plots, and leverage diagnostics. These diagnostics ensure that statistical inferences are not driven by violations of linearity, normality, or homoscedasticity.

Growth dynamics were explored using the *Orange* dataset through longitudinal visualization of tree circumference against age. Separate growth curves were plotted for each tree, allowing comparison of growth rates and patterns. This approach mirrors crop growth monitoring techniques used in precision agriculture, where repeated measurements over time inform management decisions such as irrigation, fertilization, or thinning.

Physiological response analysis was demonstrated using the *CO₂* dataset. Carbon dioxide uptake was examined as a function of ambient CO₂ concentration, plant type, and treatment condition. Grouped line plots were used to show differential response curves, highlighting how environmental stressors and plant characteristics influence productivity. This form of analysis is directly relevant to understanding crop responses to climate variability and elevated atmospheric CO₂ levels.

Experimental treatment effects were evaluated using the

PlantGrowth dataset. A one-way ANOVA was conducted to test whether mean plant weights differed across treatment groups. When the overall F-test indicated statistically significant differences, Tukey's Honest Significant Difference (HSD) post-hoc test was applied to identify which specific treatment pairs differed. This procedure reflects standard practice in agronomic field trials where multiple treatments are compared under controlled conditions.

Interpretation within a food security framework

Analytical results were interpreted through the lens of the four dimensions of food security: availability, access, utilization, and stability. Rather than focusing solely on statistical significance, emphasis was placed on substantive interpretation and policy relevance. Regression coefficients, growth trajectories, and treatment effects were linked to practical agricultural questions such as labor allocation, crop growth potential, stress tolerance, and yield enhancement.

The illustrative nature of the analyses allows the methodology to serve as a template for applied agricultural analytics. Researchers and practitioners can substitute the built-in datasets with context-specific data, such as crop yield panels, household surveys, remote sensing imagery, or sensor data, while maintaining the same analytical logic.

Ethical considerations and reproducibility

All data used in this study are publicly available and non-identifiable. As such, no ethical approval was required. Reproducibility was prioritized by relying on standard R functions and widely available datasets. This approach aligns with best practices in open science and supports capacity building in low- and middle-income countries, where access to proprietary data and software may be limited.

Results and Discussion: Illustration of analyses using built-in R datasets

It should be noted that the analyses use standard datasets in R software (swiss, CO₂, Orange and plantgrowth) to demonstrate common methods. These can be replaced or extended with domain specific datasets for example crop yield panels, remote sensing rasters or mobile data for applied projects (R Core Team, 2024).

Example A: Socioeconomic Predictors of Agricultural Labor (dataset: swiss)

In this section cross-section socio-economic data is used to explore relationships between agricultural labor share and proxies of development. This mimics how policymakers might use national household or district surveys to target agricultural support. For this study Agriculture is taken as a dependent variable. It is thought that engagement in labour as is required in agriculture is affected jointly by a number of independent variables such as education, the percentage score in examination and perhaps state of health as may be indicated by infant mortality. Modeling of agriculture is therefore developed as a function of these three variables giving a multiple regression of the form:

$$Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \epsilon \quad (1)$$

where

Y is the dependent variable representing percentage of males working in agriculture

X₁ is percentage of draftees who had the highest educational achievement (primary school completion or higher)

X₃ is the percentage of draftees (young men) who received the highest grade in army physical examinations. This variable can be used as a proxy for overall health, physical condition, and quality of human capital in a district.

X₂ is the number of infants who died per 100 live births. Infant mortality was a major demographic challenge in the 19th century in Switzerland. High mortality tended to occur in poorer, rural districts with limited access to midwives or medical services

β₀ is the constant term representing the outcome in agriculture if the independent variables had no effect

β_i = 1,2,3 represent the regression coefficients

ε the error term

The fitted model evaluates whether levels of male education, examination performance, and infant mortality are associated with the percentage of males involved in agriculture across the 47 French-speaking Swiss provinces. When the model is run in R the output given in table 1 is obtained

In table 1 the intercept has an estimated value of 104.40, with p < .001. β₀, which represents the expected value of Agriculture when the Education, Examination and Infant mortality are all equal to zero, is highly significant. The first independent variable of Education has the coefficient estimated at β₁ = -0.7513 and p = 0.0327. It is negative and statistically significant (p < 0.05). In other words every

Table 1: Regression coefficients

Coefficients:	Estimate	Std. Error	t value	Pr(> t)		
(Intercept)	104.4029	17.5451	5.951	4.31E-07	***	
Education	-0.7513	0.3405	-2.207	0.03272	*	
Examination	-1.3698	0.411	-3.333	0.00177	**	
Infant.Mortality	-1.1487	0.8097	-1.419	0.16319		

NB

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

increase of one unit in education which is the proportion of draftees with highest education results in a decrease of the percentage employed in agriculture holding all other predictors constant. One can therefore conclude that the provinces with higher educational attainment tended to have lower dependence on agriculture, consistent with structural economic transition. Education is thus a significant socio-economic predictor.

For the examination variable the coefficient is $\beta_2 = -1.3698$, and $p = 0.00177$. The coefficient is negative and highly significant ($p < 0.01$). This implies that for every unit increase in examination score, agricultural employment decreases by approximately 1.37 percentage points, when other variables are controlled. In context the provinces with stronger academic performance had even lower agricultural participation. This strengthens the interpretation that human capital development correlates with shifting away from agriculture. For the third variable, the coefficient of Infant Mortality $\beta_3 = -1.1487$, and $p = 0.1632$. This coefficient is negative and not statistically significant ($p > 0.05$). This means that there is no statistically reliable evidence that infant mortality predicts variation in agricultural employment after controlling for Education and Examination. This is how statistics can be used to enhance decision making or making relevant formulation of policies.

In the next section the model diagnostics are given with plotted graphs that can help visualizing the residuals. Fig.1. In the graph of Residuals vs Fitted, show mild




Figure 1: Model diagnostics

nonlinearity as the residuals scatter fairly randomly. For the Q-Q plot, the points are mostly on the line with a few tail deviations which show mild outliers. The residuals are close to normal and therefore acceptable. As for the Scale-Location plot the red line is almost horizontal with a mild increase in spread showing a slight heteroscedasticity. The last graph is supposed to show influential observations. Any point near or past Cook's D lines could influence

model. In the swiss data only about two are near the line but are not extreme.

Example B: Orange Tree Growth Curves (dataset: Orange)

We use the orange dataset to explore the tree growth of oranges in terms of circumference increase of their individual trees against their ages, expected yields as well as the growth anomalies that may have affected the trees under similar conditions. Fig. 2 gives the graph of the dataset of the different tree growth rates.

From Fig. 2, the growth over time depicts an upward




Figure 2: Orange tree Growth Curves


trend for all the five trees. It means circumference increases as age increases, as would always be expected biologically. The graph shows typical sigmoidal growth, where trees grow rapidly initially and then slow down as they mature. The curves do not overlap so much, meaning the trees grow at different rates. Tree 2 & Tree 4, for instance, tend to have higher circumference values at most ages suggesting they grow faster or become thicker over a shorter time. Tree 1 & Tree 3 remain comparatively lower indicating slower growth. It should be noted that since trees were measured at the same ages, divergence in curves reflects biological variability, not measurement timing.

Clearly this dataset demonstrates how data analytics helps understand growth performance. It can enhance identification of fast-growing trees, selection of superior varieties, monitoring of expected yields and detection of anomalies in growth under similar conditions for possible intervention.

Example C: Modeling plant response (dataset: CO₂)

This dataset illustrates photosynthetic uptake under varying concentrations for different plant types. They give two types of plants of Quebec and Mississippi and two treatments chilled and non-chilled plant treatments. The graphs are as shown in Fig. 3


The graphs in Fig. 3 are indicative of the fact that the



rising CO_2 concentration leads to increased uptake (a classic saturation type of response). This demonstrates the expected physiological response where photosynthesis increases as more CO_2 is available. Quebec plants tend to display higher CO_2 uptake rates compared to Mississippi plants, especially at higher concentrations. This may be reflective of the local adaptations to cooler climates, leading to more efficient carbon assimilation. It is also evident that chilled plants often show lower uptake than non-chilled plants within each type. This indicates that cold treatment suppresses physiological performance and hence ends up reducing the plant's capacity to assimilate carbon. The curves are also rising upwards steeply at the beginning, and then gradually flatten, showing that photosynthesis approaches a maximum rate (V_{max}), consistent with Michaelis-Menten kinetics in plant physiology.

Example D: Treatment Effects on yields (The dataset: PlantGrowth)

This example presents a typical case of dataset that can be studied in basic experimental design and analysis



of variance (ANOVA). It contains dry plant weights for three treatment groups namely ctrl, trt1, and trt2. Though the data is small, it is useful for illustrating yield changes under different treatment conditions. This has been used to illustrate one-way ANOVA for comparing means across groups. Fig. 4 shows the box-plot of the mean plant weight by treatment groups. Table 2 shows that treatment effect is statistically significant on the yield

Table 2: One-way ANOVA to test if mean weights differ by group

Source	Df	S u m Sq	Mean Sq	F value	Pr(>F)
group	2	3.766	1.8832	4.846	0.0159 *
Residuals	27	10.492	0.3886		

NB: Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

with $p = 0.0159 < 0.05$. Further analysis was therefore necessary to determine which of the three treatments is significant. For this we conducted Turkey's post-hoc test of multiple comparisons of means taking 95% family-wise confidence level. The results are in Table 3. From the output of the results presented in table 3, only one

Table 3: Turkey's post-hoc test of multiple comparisons of means

Group	diff	lwr	upr	p adj
trt1-ctrl	-0.371	-1.0622161	0.3202161	0.3908711
trt2-ctrl	0.494	-0.1972161	1.1852161	0.1979960
trt2-trt1	0.865	0.1737839	1.5562161	0.0120064

pair (trt2-trt1) is significantly different while the other two pairs are not. It can be concluded therefore that the plants in trt2 are significantly heavier than in trt1

Conclusion and Recommendation

From the analyses done, descriptive and regression analyses in the first example of swiss dataset reveal socio-economic correlates of agricultural reliance and can inform where complementary development investments are needed to boost productivity and access to markets. The orange dataset demonstrate growth curves that show temporal heterogeneity across trees. Clustering trees by growth patterns can help to identify groups needing similar management practices such as irrigation or pruning regimes. This idea of clustering can also be extended to farmers who can be grouped by field productivity for precision recommendations. As an analytic takeaway for agriculture, the curve analysis and unit clustering are practical for site specific management and for designing differentiated advisory services for targeted, cost-effective interventions for policy making. In the CO_2 dataset, the plot captures a fundamental ecological insight. As indicated earlier the plant type and environmental stress strongly influence CO_2

assimilation rates, which have implications for agricultural productivity and climate-resilient crop selection. Quebec plants' superior uptake suggests they may be better suited for cooler climates or for high-CO₂ environments. Lastly the plantgrowth dataset can guide extensively on the most promising growth-enhancing treatment on one hand while also showing that of inhibitory effect. This type of result is important in agricultural productivity experiments where decision on the kind of treatments that can increase biomass yield.

Bridging analytics and food security in practice From insight to action: operational challenges

Despite promising analytics, practical translation faces barriers. They include; data availability and quality where it is true that smallholder systems often lack digitized records; remote sensing may have difficulty in fragmented landscapes. Infrastructure and connectivity is another case in that rural broadband and electricity remain constraints in many regions, limiting real-time data flows and IoT deployments. This problem has been documented by many World Bank reports. Further to the challenges reported, capacity and extension is a hindrance to farmers and local extension agents who still need training to interpret and act on analytics. Interfaces must be designed for usability and linguistic or cultural fit. There is also the governance and data rights which include unclear ownership, privacy, and commercial interests. This can inhibit data sharing critical for public good analytics. Wolfert *et al.* (2017) outline data governance concerns.

Equity and inclusion

Analytics can widen inequalities if adoption favors larger, better-connected farms. Deliberate policies that include subsidized access, cooperative models, and public sector platforms are necessary to ensure smallholder inclusion. Information need to be packaged to be all-inclusive and suitable, to a great extent for all. It should be readily available and suitable for those with hearing or visual impairments as well as those of varied socio-economic levels

Advancing food security quickly

If nations have to be food secure, they must invest in technology that can enable early warning and targeted interventions. Combining satellite indices with socio-economic data supports rapid targeting of emergency inputs and social protection, reducing acute food insecurity. Post-harvest loss reduction must also be embraced. Analytics for cold chain management and logistics optimization reduce losses and increase effective food availability. Another way is adopting climate smart planning which entails integrating climate projections with cropping systems models. This supports adaptation and insurance design usually referred to as climate smart agriculture

Generally understanding these challenges and of course making deliberate effort to counter them while

acknowledging the crucial role of data analytics in agriculture helps in evaluation and choice of better pathways that can boost productivity and strengthen food security.

Policy recommendations and implementation roadmap

1. Develop national digital agriculture roadmaps. Governments should set interoperability standards, data governance rules, and fund public good data platforms to lower barriers for innovators (World Bank playbook).
2. Invest in rural connectivity and sensor networks. Public-private partnerships can extend connectivity and subsidize shared sensor services (soil probes, weather stations).
3. Build data literacy and extension capacity. Train extension workers in data-driven advisories and co-design tools with farmers to ensure usability.
4. Support open data and platforms for smallholders. Public datasets such as weather, soil maps, and satellite products should be accessible; encourage interoperable private platforms that can plug into public services.
5. Pilot blended finance for scaling analytics. Use concessional finance and results-based instruments to de-risk investments in digital extension and precision tech for smallholder contexts.
6. Prioritize applications with strong food-security payoffs. Early warning systems, post-harvest logistics, and targeted input delivery often yield high social returns and should be prioritized.

CONCLUSION

Data analytics is not a panacea, but when combined with sound policy, investments in connectivity and capacity, and attention to equity, it can be a powerful lever to raise agricultural productivity and strengthen food security. Translating analytics into inclusive outcomes requires bridging technical innovation with institutional reform and ground level engagement with farmers.

REFERENCES

- Aissi, M. E., Benjelloun, S., Lakhrissi, Y., & Ali, S. E. (2023). A Scalable Smart Farming Big Data Platform for Real-Time and Batch Processing Based on Lambda Architecture . *Journal of System and Management Sciences*, 13 (2), 17-30.
- Dogho, M. O., & Babatunde, I. O. (2025). Data Analytics in Food Safety: Improving Quality Control and Preventing Contamination . *Current Journal of Applied Science and Technology*, 44 (4), 245-256.
- FAO, IFAD, UNICEF, WFP & WHO. (2025). Annual Report. The State of Food Security and Nutrition in the World (SOFI). Food and Agriculture Organization of the United Nations .
- Lohit, V. S., & Mujahid, M. M. (2022). Big Data Analytics in Developing Smart and Sustainable Solutions for the Agricultural Industry 1 . *Technoarete Transactions on Advances in Data Science and Analytics*, 1 (2), 1-6.

- R Core Team. (2024). A language and environment for statistical computing. R Foundation for Statistical Computing.
- Sarker, M. R., Abdolrasol, M. G., Md, S. M., Kadir, R. A., Ahmad, M. N., & Olazagoitia, J. L. (2025, September). Advancing Agriculture Automation Systems: Technological Innovations, Possible Applications, Challenges, and Recommendations. *Advances in Agriculture* , 1-33.
- Soussi, A., Zero, E., Sacile, R., Trinchero, D., & Fossa, M. (2025). Smart Sensors and Smart Data for Precision Agriculture: A Review. *Sensors* , 1-32.
- Wang, Z., Akber, M. A., & Aziz, A. A. (2025, August 14). Application of Remote Sensing and Geographic Information Systems for Monitoring and Managing Chili Crops: A Systematic Review. *Remote Sensing* , 1-23.
- Wolfert, S., Ge, L., Verdouw, C., & Bogaardt, M.-J. (2017). Big Data in Smart Farming – A review. *Agricultural Systems*, 153, 69-80.
- World Bank. (2025). Digital Agriculture Roadplaybook and technical guidance for national digital agriculture strategies.
- Yang, Z., Wu, W., Di, L., & Üstündağ, B. (2017). Remote sensing for agricultural applications. *Journal of Integrative Agriculture* 2017, 16(2): 239–241, 16 (2), 239-241.
- Yu, L., Du, Z. L., Zheng, J., Zhao, Q., Wu, H., Weise, D., et al. (2025). Enhancing global agricultural monitoring system for climate-smart agriculture. *Climate Smart Agriculture*, 2, 1-10.