



DSPACE

<https://dspace.org/>

Effective Dose of Crude Ethanol and Aqueous Extracts of Phytolacca Dodecandra (L' Herit) on Anopheles Gambiae (Diptera: Culicidae) Larvae

Owiti, Yugi Jared

2015

Journal of Mosquito Research 2015, Vol.5, No.11, 1-6

<http://ir-library.kabianga.ac.ke/handle/123456789/660>

Yugi J.O., Okeyo-Owour, J B., Were-Kogogo P., Luvaha G.M., Okwiri L.O., Rono T.M and Vulule M.J., 2015, Effective Dose of Crude Ethanol and Aqueous Extracts of Phytolacca Dodecandra (L' Herit) on Anopheles Gambiae (Diptera: Culicidae) Larvae, Journal of Mosquito Research, Vol.5, No.11 1-6 (doi

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/282962563>

Effective Dose of Crude Ethanol and Aqueous Extracts of *Phytolacca Dodecandra* (L' Herit) on *Anopheles Gambiae* (Diptera: Culicidae) Larvae

Article · January 2015

DOI: 10.5376/jjmr.2015.05.0011

CITATION

1

READS

58

1 author:



Jared Owiti Yugi

University of Kabianga

6 PUBLICATIONS 8 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Impact of geophagy on health and work output of gravid tea pickers [View project](#)

Effective Dose of Crude Ethanol and Aqueous Extracts of *Phytolacca Dodecandra* (L. Herit) on *Anopheles Gambiae* (Diptera: Culicidae) Larvae

Yugi J.O.^{1,€}, Okeyo Owour J.B.², Were Kogogo P.³, Luvaha G.M.¹, Okwiri L.O.¹, Rono T.M.⁴, Vulule M.J.⁵

1. Department of Applied Sciences, Kisumu Polytechnic, P. O. Box 40100, Kisumu, Kenya

2. School of Agriculture, Natural resource and Environmental Studies, Rongo University College, Box 103404, Rong, Kenya

3. School of Biological and Physical Sciences, Jaramogi Oginga Odinga University, P. O. Box 20000, Rongo, Kenya

4. University of Eldoret, P. O. Box 1125100, Eldoret, Kenya

5. Kenya Medical research Institute, Centre for Global Health Research, P. O. Box 19280, Kisumu Kenya

€ current address: Department of Biological Sciences, University of Kabianga, P. O. Box 20200, Kericho, Kenya

Corresponding author email: yugijared@gmail.com

Journal of Mosquito Research, 2015, Vol.5, No.11 doi: 10.5376/jmr.2015.05.0011

Received 17 Jun, 2015

Accepted 18 Jul, 2015

Published 04 Aug., 2015

Copyright © 2015 Yugi et al, This is an open access article published under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Preferred citation for this article

Yugi J.O., Okeyo Owour, J.B., Were Kogogo P., Luvaha G.M., Okwiri L.O., Rono T.M and Vulule M.J. 2015, Effective Dose of Crude Ethanol and Aqueous Extracts of *Phytolacca Dodecandra* (L. Herit) on *Anopheles Gambiae* (Diptera: Culicidae) Larvae. *Journal of Mosquito Research* Vol.5, No.11 1-6 (doi: [10.5376/jmr.2015.05.0011](http://dx.doi.org/10.5376/jmr.2015.05.0011))

Abstract Objective: malaria vector control is either by adult or larvae management. This study reports on the effective crude ethanol and aqueous extracts of leaf and mature green fruits of *Phytolacca dodecandra* (L. Herit) as a larvicide against *Anopheles gambiae* (Diptera: Culicidae) mosquitoes.

Material & Methods Batches of twenty freshly hatched or moulted larvae (1st, 2nd, 3rd and 4th instars) of *An. gambiae* were transferred using droppers into plastic containers measuring 6 cm mouth and 5.7 cm base diameter by 3.5 cm height arranged in sets. Each container contained an approximate of 33 millilitre solution of a particular concentration of leaf (shoot or midsection) or fruit extracts of *Endod*. *Endod* plants were sourced from Nyando and Eldoret. The solutions were a serial dilution of 40, 20, 10, 5 and 5mg/100mls of the crude extracts. Solutions of *Neem* and *deltamethrin* were used as positive control and that of water only as negative control. Three replicates were set up for each concentration. WHO threshold of > 80 % mortality was used as standard to assess effectiveness of the larvicide.

Results Over 80% of all larvae (L1s, L2s, L3s and L4s) died from exposure to concentrations of 20mg and higher of ethanol extracts of mature green fruits of *Endod* irrespective of source. Only 40mg/100mls of aqueous extracts of *Endod* leaves of the shoot sourced from Eldoret killed > 80% of exposed L1s. Higher mortalities were observed for extracts from leaves of the shoot of *Endod* than those from the midsection.

Conclusion: Higher doses of ethanol and water extracts of *Endod* leaves and fruits were lethal to all *Anopheles gambiae* larvae.

Keywords *Anopheles gambiae* larvae, *Phytolacca dodecandra*, *Neem*, *Deltamethrin*

Introduction

Synthetic insecticides (Poopathi and Archana, 2010) have for a long time been used to manage extensive morbidity and mortality within disease endemic countries (Boutayeb, 2006) due to Mosquito-borne diseases. Though effective this strategy is accompanied with serious harm to human and other animals in the environment (Cartilla and De la Cruz, 2012). In addition continued use of synthetic insecticides has been demonstrated to lead to reduced control due to development of resistant in insect populations (Charles and Nielser LeRoux, 2000).

The increasing negative effect of synthetic insecticides has made scientists seek for alternative friendly pest control methods (Poopathi and Archana, 2010; Montasser et al., 2011). Botanicals such as *Phytolacca dodecandra* here after *Endod* have been found promising as alternative larvicidals against *Culex quinquefasciatus* (Misganaw et al., 2012) and *An. gambiae* (Yugi et al., 2015) and pupicidals against *Culex quinquefasciatus* (Misganaw et al., 2012). There is also evidence of demonstrable microbial activity (Joshi et al., 2011) Though efficacy of *Endod* extracts on *An. gambiae* larvae has been demonstrated, no information exists on the dose or amount of extracts

that are effective on each larval stage. It is in line with 2 Discussions

this that this study was designed to study and report Ethanol extracts of mature green fruits of Endod at on the effective dose of crude ethanol and water higher concentrations killed exposed aquatic stages extracts of mature green fruits and leaves of Endod of An. gambiae Mortality percentages however, larvae of An. gambiae in a laboratory set up. reduced with continued dilution. This trend was similar to Ethanol extracts of Endod leaves although

1 Results

The experiment was conducted for a period of eight months using a total of 84,240 larvae (L1s, L2s, L3s and L4s). WHO threshold of >80% mortality was used as standard to decide on the effectiveness of the various doses against the exposed larvae. L1s, concentration of 20 mgs and 40 mgs of ethanol extracts of mature green fruits of Endod from Eldoret and Nyando respectively killed over 80% of the exposed larvae. Earlier the same trend had been reported for powdered extracts of Endod on An. gambiae larvae (Were, 2008). This demonstrates that extracts of ethanol and water extracts of Endod have similar efficacies as Chenopodium ambrosioides (Jude et al., 2013) and Jatropha curcas (Zewdneh et al., 2011) against An. gambiae and Anopheles arabiensis larvae respectively.

For aqueous extracts of Endod mature green fruits and leaves of the shoot, mortality of over 80% was observed for concentrations of 40mgs only irrespective of source (Figure 1). Extracts from both mature green fruits and leaves of Endod were potent against An. gambiae larvae though potency of leaf extracts were lower than that of fruits. Potency of the extracts also differed with the vertical

position of leaves on the Endod plant such that ethanol extracts of leaves found at the shoot were more potent than those found at the midsection of the plant. This showed that the probable distribution of the bioactive on Endod plant are such that fruits > leaves at the shoot > leaves at the midsection.

For L2s, concentrations of 20 mg and 40 mg of ethanol extracts of mature green fruits from Eldoret and Nyando killed over 80% of exposed L1s respectively. Mortality of over 80% was observed for concentrations of 40 mgs of extracts of leaves of the shoot and midsection of Endod from Nyando and Eldoret respectively. Concentrations of 20 mg and 40 mgs of aqueous extracts of mature green fruits from Eldoret and Nyando killed over 80% of exposed L2s respectively (Figure 2). Ethanol extracts of Endod parts were more toxic than aqueous extracts of the same parts for all concentrations against An. gambiae larvae. Solvent type used in extraction had been demonstrated to determine

potency of extracted bioactives (Anupam et al., 2012) as well as larviciding potency. This had been demonstrated for Solanum xanthocarpum (Mohan et al., 2006), Euphorbia tirucalli (Singh et al., 2007), Eucalyptus globules (Maurya et al., 2007), Citrullus colocynthis (Sakthivadivel and Daniel, 2008), Azadirachta indica (Mgbemena, 2010) and Solanum nigrum (Raghavendra et al., 2009).

For L3s, concentrations of 10 mg and 20 mg and above of ethanol extracts of mature green fruits and leaves of the shoot of Endod from Eldoret and Nyando respectively killed over 80% of exposed L3s. The same was observed for concentration of 40 mgs of ethanol extracts of leaves of the midsection of Endod (Figure 3). For the L4s, more than 80% mortality of exposed larvae was observed for concentrations of 10mgs and above for ethanol extracts of mature green fruits irrespective of source (Figure 4). Ethanol extracts of Endod parts sourced from Eldoret were more potent than of the same parts sourced from Nyando. This showed that geographical origin of the plants played a role in the concentration and

distribution of bioactives within the plants (Anupam et al., 2012). A similar observation had been made on Endod parts (Were, 2008), Citrus sp, Ocimum sanctum and Azadirachta indica (Mgbemena, 2010) and Jatropha sp (Sakthivadivel and Daniel, 2008) just to mention but a few. Mortality of over 80% was observed for L1s and L3s for ethanol and aqueous extracts respectively for Neem while Deltamethrin killed all exposed larvae irrespective of stage or concentration. The effects of the extracts were considered significantly different at $p < 0.05$ (Table 1) for all exposed larval stages

Figure 1 Mortality of *An. gambiae* first instars (L1) exposed to different concentrations of crude ethanol and water extract of *Endod.* Error bars represent standard error of means

Figure 2 Mortality of *An. gambiae* second instars (L2) exposed to different concentrations of crude ethanol and water *Endod* extract. Error bars represent standard error of means

This study demonstrates that crude ethanol and water extracts of *Endod* are potent against *An. gambiae* larvae at high doses. It also demonstrates that solvent type used in the extraction influences activity of extracts from *Endod* irrespective of parts or geographical origin.

3 Materials and Methods

3.1 Study area, experimental mosquitoes and study design

The experiments were conducted in the Entomology

laboratory at Centre for Global research Institute/ Kenya medical research Institute (CGHR/KEMRI). Pink eyed *An. gambiae* mosquitoes maintained at the laboratories and reared following standard techniques (Das et al., 2007) were used in the experiments. A completely randomized informal, after only with control, experimental design (Kothari, 2004) was used to investigate the larvicidal effect of crude ethanol and water extracts of *Endod* on the mosquito larvae. The condition in the insectary was as described elsewhere (Yugi et al., 2014).

Figure 3 Mortality of *An. gambiae* third instars (L3) exposed to different concentrations of crude ethanol and water Endod extracts. Error bars represents standard error of means

Figure 4 Mortality of *An. gambiae* fourth instars (L4s) exposed to different concentrations of crude ethanol and water Endod extract. Error bars represents standard error of means

Table 1 Duncan's statistics on mortalities due to exposure of *An. gambiae* larvae to crude extracts of parts of Endod

Stage	df	Treatments							Neem	Deltamethrin
		Parts of Endod used								
		Eldoret			Nyando					
		Fruits	Shoot	Midsection	Fruits	Shoot	Midsection			
L1	4	0.0128	0.0001	0.0001	0.0145	0.1037	0.3729	0.0001	-	
L2	4	0.0449	0.0229	0.0015	0.0660	0.0296	0.0001	0.0008	-	
L3	4	0.0603	0.0234	0.2632	0.0399	0.0830	0.0490	0.0001	-	
L4	4	0.1679	0.2245	0.0401	0.0037	0.0867	0.0356	0.9642	-	

Note: 1 P is the probability for the level of significance. P was considered significant at $p < 0.05$

2 df is the degree of freedom of the considered units

3.2 Deltamethrin (KOTab 1-2-3®) and plant materials acquisition and preparation

Deltamethrin (KOTab 1-2-3®) was acquired and prepared as described (Yugi et al., 2015). Fresh leaves (shoot and midsection) and mature green fruits of Endod and Fresh leaves of *Azadirachta indica* (Neem), were acquired, identified and voucher specimen deposited as described elsewhere (Yugi et al., 2015). The plant parts were dried in a shade at room temperature, grounded and extracts obtained using ethanol and water as described in details elsewhere (Tilahun et al., 2003; Parekhet al., 2005; Das et al., 2010; Yugi et al., 2015).

The extracts were concentrated by freeze drying using a rotary vacuum evaporator at -40°C to obtain essential oil that was then kept in airtight glass bottles to serve as stock quantity. From the freeze dried stock, 80mg were weighed and serial dilutions made to obtain different concentrations of 40, 20, 10, 10, 5 and 2.5 mg in 100mls of rain harvested water.

3.3 Larvicidal bioassays

Larvicidal activities were tested in accordance with the WHO procedure (WHO, 1996) and as described elsewhere (Yugi et al., 2015). Mortality rate were registered after 24 hour exposure period and larval mortality calculated for each concentration using the formula;

Standard WHO procedures were used to assess effectiveness of the extracts as larvicide at a mortality rate of > 80% (WHO, 2005). Moribund and dead larvae were collected and disposed off in a septic tank.

3.4 Data analysis

Data obtained from the bioassays was entered in excel spreadsheet for ease of handling. The relationship between the effective doses of the extracts with respect to parts of Endod used was determined using descriptive statistics. One way analysis of variance (ANOVA) was used to assess the level of significance of the various doses on larvae mortality. All statistical analysis was performed using SAS statistical package version 20.

Competing interest

The authors declare that they have no competing interest.

Acknowledgements

We thank Patience Akoth, Charlotte Awuor, Hilda Ochieng, Trevor Omondi and Richard Amoto, for helping culture all mosquitoes used in this study. We thank Kisumu Polytechnic for equipments for the extractions of the crude extracts from Endod and Neem plants, Centre for Global Health Research/Kenya Medical Research Institute (CGHR/KEMRI) for mosquitoes, laboratory space and equipments, VIRED International for providing transportation and logistics for sourcing for Endod and Neem parts. We also thank National Commission for Science, Technology and Innovation (NACOSTI), Kenya for funding this project.

References

- Anupam G., Nandita C., and Goutam C., 2012, Plant extracts as potential mosquito larvicides. *Indian Journal of Medical Research*, 135:5981
- Boutayeb, A., 2006, The Double burden of communicable and noncommunicable diseases in developing countries. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 100(3): 191-199
<http://dx.doi.org/10.1016/j.trstmh.2005.07.021>
- Cartilla P., and De la Cruz J., 2012, Termiticidal Potential of *Schytarpheta Jamaicensis* (L.) Yahl, 1: 1-5
- Charles C., and Nielsen Roux C., 2000, Mosquitocidal Bacterial Toxins: Diversity, Mode of Action and Resistance. *Phenomena*, 95(Suppl 1): 201-206
<http://dx.doi.org/10.1590/s0070-762000000700034>
- Das K., Tiwari R.K.S., and Shrivastava D.K., 2010, Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends. *Journal of Medicinal Plants Research*, 4(2): 104-111
- Das N.G., Goswami D., and Rabha B., 2007, Preliminary evaluation of mosquito larvicidal efficacy of plant extracts. *Journal of Vector Borne Diseases* 44: 145-148
- Joshi B., Sah G.P., Basnet B.B., Bhatt M.R., Sharma D., Subedi K., Pandey J., and Malla R., 2011, Phytochemical extraction and antimicrobial properties of different medicinal plants: *Scimunc sanctum* (Tulsi), *Eugenia caryophyllata* (Clove), *Achyranthes bidentata* (Dativan) and *Azadirachta indica* (Neem). *Journal of Microbiology and Antimicrobials* 1(1): 1-7
- Jude D.B., Petola A.S., Sali A.N., Judith L.N., Maximilienne N., Julius E.O., and Rose G.F.L., 2018, Larvicidal and Repellent Potential of *Chenopodium ambrosioides* Linn Essential Oil against *Anopheles gambiae* Giles (Diptera: Culicidae). *The Open Entomology Journal*, 16-22
<http://dx.doi.org/10.2174/1874407901307010016>
- Kothari C.R., 2004, Research design: research methodology, methods and techniques. 2nd edition. New Age International Publishers, New Delhi, India, 4750
- Maurya P., Mohan L., Sharma P., Batabyal L., and Srivastava C.N., 2007, Larvicidal efficacy of *Aloe barbadensis* and *Cannabis sativa* against the malaria vector *Anopheles stephensi* (Diptera: Culicidae). *Entomological Reserves* 37: 153-156
<http://dx.doi.org/10.1111/j.1745967.2007.00105.x>
- Mgbemena I.C., 2010, Comparative evaluation of larvicidal potentials of three plant extracts of *Aedes aegypti*. *Journal of American Science* 6: 435-440
- Misganaw N., Moges S., Tadele M., Tesera M., Temesgen T., and Raja N., 2012, Evaluation of Multi Potential Bioactive Endod *Phytolacca dodecandra* (L. Herit) Berries Extracts Against Immature Filariid Vector *Culex quinquefasciatus* Say (Diptera: Culicidae). *Research Journal of Environmental and Earth Sciences*, 4(7): 697-703
- Mohan L., Sharma P., and Shrivastava C.N., 2006, Evaluation of *Solanum xanthocarpum* extracts as a synergist for cypermethrin against larvae of filarial vector *Culex quinquefasciatus* (Say). *Entomological Reserves* 36: 220-225
<http://dx.doi.org/10.1111/j.1745967.2006.00037.x>

- Montasser S.A., Shamseldean M.M., Mahmoud Y.A., Ibrahim S.A., and Tilahun G., Egualé T., and Kassa 2003, Effect of Endod on parasitic leech, *Lymanis nilotica* Bulletin of Animal Health and Production 51: 75-81
- Sharaby A., 2011, Plant Extracts as Alternative Botanical Insecticides for Control the Grasshopper *Heteracris littoralis* Ramb. (Orthoptera: Acrididae) with Reference to Histological Changes the Reproductive System. Journal of Basic Application of Science Resources, 1(9): 1031-1038
- Parekh J., Jadeja D., and Chanda S., 2005, Efficacy of aqueous and methanol extracts of some medicinal plants for potential antibacterial activity. Turkish Journal of Biology, 29: 202-210
- Poopathi S., and Archana B., 2010, Mosquitocidal bacterial to *Bacillus sphaericus* and *Bacillus thuringiensis* serovar israelensis. Mode of action, cytopathological effects and mechanism of resistance, 50(1): 65
- Raghavendra K., Singh S.P., Subbarao S.K., and Dash A.P., 2009, Laboratory studies on mosquito larvicidal efficacy of aqueous & hexane extracts of dried fruit of *Solanum nigrum*. Indian Journal of Medicinal Research, 130: 74-77
- Sakthivadivel M., and Daniel T., 2008, Evaluation of certain insecticidal plants for the control of vector mosquitoes *Culex quinquefasciatus* and *Anopheles stephensi* and *Aedes aegypti*. Application of Entomological Zoology, 43: 57-63
<http://dx.doi.org/10.1303/aez.2008.57>
- Singh R.K., Dhiman R.C., Mittal P.K., 2007, Studies on mosquito larvicidal properties of *Eucalyptus citriodora* Hook (family Myrtaceae). Journal of Communicable Diseases, 39: 233-236
- WHO 1996, Report of the WHO informal Consultation on Evaluation and Testing of Insecticides. Geneva, 37
- WHO 2005, Guidelines for laboratory and field testing of mosquito larvicides. World health organization communicable disease control, prevention and eradication who pesticide evaluation scheme. WHO/CDS/WHOPES/GCDPP/2005, 13
- Yugi J.O., Okeye Owuor J.B., Atieli F., Amato R., and Vulule J.M., 2014, Knockdown effect of crude ethanol extracts of *Phytolacca dodecandra* on *Anopheles gambiae* adults. Journal of Mosquito Research, 4(18): 17
<http://dx.doi.org/10.5376/jmr.2014.04.0018>
- Yugi J.O., Okeye Owuor J.B., Auma C.A., Juma J.I., and Vulule J.M., 2015, Larviciding potency of water and ethanol extracts of *Phytolacca dodecandra* (L. Herit) on *Anopheles gambiae* (Diptera: Culicidae). Journal of Mosquito Research, 5(2): 1-6
- Zewdneh T., Mamuye H., Asegid T., Yalemtehay M., Bayene P., 2011, Larvicidal effects of *Jatropha curcas* L. against *Anopheles arabiensis* (Diptera: Culicidae). CNCS, Mekelle University 3(1):52-64