



DSPACE

<https://dspace.org/>

Effects of Climate Variability on the Outbreak and Spread of Newcastle Disease in Suneka Division of Kisii County

Nyaiyo, Norah M.; Maangi, Eric Nyankanga

2016

Research on Humanities and Social Sciences

<http://ir-library.kabianga.ac.ke/handle/123456789/661>

Nyaiyo, N. M., & Maangi, E. N. Effects of Climate Variability on the Outbreak and Spread of Newcastle Disease in Suneka Division of Kisii County.

Effects of Climate Variability on the Outbreak and Spread of Newcastle Disease in Suneka Division of Kisii County

Norah M.Nyaiyo^{1*} and Dr.Eric Nyankanga Maangi²

1. Doctoral student and part-time Lecturer, Kisii University, Kenya

2. Lecturer, Department of Psychology and Foundations University of Kabianga, Kenya

*corresponding author

ABSTRACT

Newcastle Disease (ND) is a common occurrence in Suneka Division. ND is caused by Avian Paramyxovirus Type 1, is one of the most significant diseases for poultry producers around the world. This poultry disease is influenced directly or indirectly by weather as it affects the timing and the intensity of an outbreak. This article examined the influence rainfall and temperature on the outbreak and the spread of Newcastle disease in Suneka Division of Kisii County. The study used a descriptive design with both qualitative and quantitative approaches. The sample size was 200 households and respondents were poultry farmers. Veterinary officers were key informants. Primary data was collected using questionnaires and interview schedules. Secondary data on rainfall and temperature was collected from the Kisii meteorological station for 2007 and 2011. Data was analysed using measures of central tendency such as means and mode. They were presented using frequency tables. Karl Pearson coefficient of correlation was used to test the null hypothesis that there was no relationship between temperature and rainfall, and the occurrence of ND at 5% level of significance. The study revealed that ND occurs during the months of July-August and December-January. It was generally found out that this disease occurs after the long March, April and May (MAM) and short rains October November and December (OND). It was also found out that there was a positive Pearson Correlation between rainfall and ND in the months of August ($r = 0.286$) and December ($r = 0.275$). However, there was a negative relationship in the months July ($r = -0.549$) and January ($r = -0.144$). Concerning temperature, there was a positive Pearson Correlation between temperature and ND in August ($r = 0.772$), July ($r = 0.683$) and January ($r = 0.159$). But, in the month of December there was a negative Pearson Correlation ($r = -0.546$). To curb the ND menace and increase productivity of free range indigenous chicken, the research recommended that there was need for farmers to monitor weather changes by obtaining data from the meteorological weather stations. Educate small-scale farmers on how to adopt hygienic and bio-safe poultry rearing methods to minimize loss of chicken through diseases, pests and predation.

Keywords: Free range, Indigenous chicken, Newcastle Disease

Introduction

Poultry plays a key role in the lives of the poor rural population in developing countries. This is especially so in Sub-Saharan Africa (SSA) where they provide income, capital assets, and fertilizers (Millar *et al.*, 2008). In Kenya both hybrid and indigenous poultry are reared. Indigenous poultry production presents a significant portion of the economy and a source of income to small scale farmers (Kaingu *et al.*, 2010). Most farmers in rural areas rear chicken because they are raised with relatively low capital, space and readily available household labour. They are also hardy, adapt well to the rural environments, survive on low inputs and adapt to fluctuations in the available feed resources (Gichoni and Maina, 1992). FRIC play a vital role in the improvement of the nutritional status and income of many poor rural households. They are also a global asset for many millions who live below the poverty line (Alders and Spradbrow, 2001).

Despite their importance, FRIC are faced by many challenges. These include: predation by snakes and birds of prey, poor housing, poor nutrition, climate variability, attack by pests and diseases and lack of adequate assistance from extension services (Moreki, 2006).

Newcastle disease (ND), caused by Avian Paramyxovirus Type 1 (APMV-1), is one of the most significant diseases for poultry producers around the world (Okwor and Eze, 2011). This poultry disease is influenced directly or indirectly by weather and climate. These may be spatial with climate affecting distribution or temporal with weather affecting timing of an outbreak and both relate to the intensity of an outbreak. Outbreaks are often associated with alternating heavy rainfall, drought and high temperatures (Perry *et al.*, 2002).

Newcastle Disease was first recognized in Indonesia in 1926 and has persisted as the major disease affecting poultry. Although little systematic research has been conducted on the epidemiology of ND in Indonesia, it is reputed to affect both indigenous and imported species of poultry throughout the country on what would appear to be a seasonal basis. Highest mortalities have been reported towards the end of the dry season for the months between July-September, the period which farmers refer to as '*musim penyakit*' or the 'disease season' (Kingston and Creswell, 1982).

A study conducted by Njagi *et al.*, (2010) revealed that Newcastle Disease Virus (NDV) was significantly higher (17.8 percent) in the dry hot zones compared to the cool wet zone at 9 percent showing climate as a risk factor in the occurrence of NDV in FRIC. Also Sonaiya (1999a) reported that the major outbreak of Newcastle disease regularly occur at the peak of the rain (June/July) and the dry season (January/February) during which mortality reaches 70 – 100% in Nigeria. Reports from Kenya (Anonymous 1996) and Ethiopia (Sonaiya 1999b) indicated that severity of this disease reached the peak in wet season but in dry season in West Africa (Mukiibi–Muka 1992; Gueye 1998). Unfortunately the period of high demand in dry season coincided with high incidence of Newcastle disease thus increasing its spread (Kuzonga *et al.*, 2008). This high demand for FRIC in December/January for Christmas and New Year Celebrations is a major factor for its spread in Nigeria. Also a study done by Olabode *et al.*, (2012) in Ilorin Kwara state in Nigeria revealed seasonal distribution of the disease indicated a higher occurrence in the dry season (October-March). Therefore pathogens or parasites that are sensitive to moist or dry conditions may be affected by changes in precipitation, soil moisture and frequent floods.

In relation to temperature, the virus can survive for more than 8 weeks in hot dry areas at temperatures of 40°C (Warner, 1989), for about 3 months at 20°C to 30°C and even longer at cooler temperatures (Lancaster, 1966). At 23°C -29°C, the virus (APMV-I) is reported to survive in contaminated litter for 10 to 14 days and 20°C in soil for 22 days (Institute of International Co-operation in Animal Biologics IICAB, 2005). When temperatures are just above freezing point (1-2°C) the virus is reported to survive on chicken skin up to 160 days and in bone marrow for nearly 200 days. APMV-1 can be inactivated by heat of 560 c for three hours or 600c for 30 minutes (IICAB, 2005).

METHODOLOGY

The study adopted a descriptive research design. The variables were ND, temperature and rainfall. Temperature is indicative of thermal stress while rainfall affects both the quality and quantity of feeds consumed. The research used simple random and purposive sampling methods. The sample size was 200 households. Both primary and secondary data were collected. Primary data was collected from poultry farmers using questionnaires. Data collected included, season of ND occurrence over 2007-2011, mortality of FRIC, socio-economic impacts of ND to poultry farmers and mitigation and adaptive measures put in place to manage ND. Data from the farmers was collected from September to December 2012. Interview schedules were used for veterinary officers. Secondary data on climatic variables (temperature, humidity and rainfall) on monthly basis was collected from Kisii Meteorological Department for 2007-2011.

Descriptive data was analysed by objectives and hypothesis. Climatic variables were analysed using the arithmetic mean for the five years. Hypothesis testing was done using the Karl Pearson Coefficient of Correlation. The Statistical Package for Social Sciences (SPSS) was used in testing of hypotheses and in the analysis of empirical data.

RESULTS AND DISCUSSIONS

Relationship between temperature and the occurrence of Newcastle Disease in FRIC in Suneka Division

Climate variability can affect poultry both directly and indirectly. Direct effects are from air temperature which influence poultry performance, growth and egg production. Indirect effects include climatic influences on severity and distribution of poultry diseases and parasites. In relation to ND, changes in temperature influenced the outbreak and severity of the disease in FRIC. When respondents were asked if changes in temperature had influence on the outbreak and spread of ND, 98.5 percent agreed that it had while 1.5 percent had no knowledge about it. This is in agreement with Dontwi *et al.*, (2011) who found out that wet season or dry season predisposes the FRIC to diseases. Further, when respondents were asked the months in which ND occurred, 74 percent reported that it occurred in July-August, 24 percent December- January and only 2 percent were not sure. Relating these months to temperature, rainfall and humidity, it was found out that temperature for July was the lowest for the five years. The temperatures recorded in the month of August were higher than July for the five years.

The mean temperature for the five years was in January was 21.8°C while for December was 21°C, July was 19.7°C while in August 20.3°C. Generally July recorded the lowest temperatures while January was among the second highest after February. The mean annual temperatures are higher for the four months. This agrees with Njagi *et al* (2010) who found out that NDV was significantly higher in drier zones.

There was a positive Pearson Correlation ($r = 0.683$) between temperature and ND in the month of July, ($r = 0.772$) in the month of August and in the month of January ($r = 0.159$). However in the month of December there was a negative Pearson ($r = -0.546$) between temperature and ND.

Table 1: Total monthly rainfall in mm for the years 2007-2011 from Kisii Meteorological Department

Year/ Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
2007	188.6	179.5	151.1	172.3	246.7	191.4	139.3	70.7	158.9	125.4	97.5	96.6
2008	30.8	59.5	304.1	291.1	118.5	136.0	220.6	191.7	168.9	241.0	154.3	82.1
2009	120.8	54.7	252.2	242.0	311.1	183.1	70.0	213.0	145.3	94.1	133.4	310.0
2010	108.6	106.5	217.7	244.6	375.7	252.3	80.3	178.1	256.7	256.9	109.3	170.3
2011	101.0	44.7	141.9	228.4	235.1	94.4	99.1	226.4	226.0	209.2	312.6	220.2
Mean	110	89	213.4	235.7	257.4	171.4	121.9	176	191.2	185.3	161.4	175.8

The mean annual rainfall for five years in January was 110 mm; July was 121mm, August 176mm and December 175.8 mm. This shows that ND occurs after the long rains MAM and short rains OND. There is a negative Pearson Correlation ($r = -0.549$ and $r = -0.144$) between rainfall and ND in the month of July and January, a positive Pearson Correlation ($r = 0.286$ and 0.275) in the month of August, and December respectively.

Conclusion

The research found that Newcastle Disease outbreaks were experienced in the months of July to August and December to January. The mean annual temperature for the five years in January was 21.8 °C while for December was 21.0°C, July was 19.7 °C while in August 20.3 °C. Generally July recorded the lowest temperatures while January was the second highest after February. There was a positive Pearson Correlation ($r = 0.772$) between temperature and ND in August, a negative correlation ($r = -0.546$) in December, a positive correlation ($r = 0.683$) in July and a positive correlation ($r = 0.159$) between temperature and ND in January.

The mean annual rainfall for five years in January was 110 mm, July was 121mm, August 176mm and December 175.8 mm. This shows that ND occurs after the long rains MAM and short rains OND. There was a negative Pearson Correlation ($r = -0.549$) between rainfall and ND in the month of July. There was a positive correlation ($r = 0.286$) in August, a positive correlation ($r = 0.275$) in December and a negative correlation ($r = -0.144$) between rainfall and ND in January.

Recommendations

On the basis of the findings of this study, the following are the recommendations that would improve the production in free range indigenous chicken, prediction and control of Newcastle disease.

- Decentralize weather stations to location levels to enable monitoring of changes in temperature, rainfall and humidity so as to predict the occurrence of this disease and take early precautionary measures.
- The government should employ more extension officers and decentralize veterinary offices to the sub-location levels so as to bring services to the farmer. This will enable farmers to get technical assistance without travelling long distances.
- Establish quarantine system to curb the spread of the disease and also establish a slaughter house to allow inspection. This will ensure that only healthy chicken are sold to consumers.
- Government to provide free drugs for vaccination and reduce the cost of feeds in order to reduce the cost of production for the poultry farmers

REFERENCES

- D.J. Alexander, Newcastle Disease, Methods of spread. In D.J. Alexander (Ed.), *Newcastle Disease*. Boston: Iowa State University Press, 1988, pp. 256-272.
- D.J. Alexander, Newcastle Disease and other Avian Paramyxovirus Infections. In, B. W. Calnek, , H. J. Burne, C. W. Beard, L.R. McDougald &, Y. M. Saif (Eds), *Diseases of Poultry* 10th ed., Iowa: Iowa State University Press, 1997.
- R. Alders and P.B. Spradbrow, Controlling Newcastle Disease in Village Chickens: A Field Manual, 2001, retrieved on 01/5/ 2012 from <http://www.ncbi.nlm.nih.gov/pubmed/10816836>
- Anonymous, Ethno veterinary medicine in Kenya: a field Manual of traditional animal health care practices. Nairobi, Kenya, 1996.
- M.A. Awan, M.J. Otte, and A.D James, The Epidemiology of Newcastle disease in rural poultry: A review, *University of Reading*: UK, 1994.
- R. Bennett, The Direct Costs of Livestock Disease: The Development of a System of Models for the Analysis of 30 Endemic Livestock Diseases in Great Britain. *Journal of Agricultural Economics* 54(1), 55-71, 2003.
- D.K. Byraru, J.L. Nakavum, and A. State, Control of Newcastle Disease in Village poultry using Thermostable ND vaccine: Presentation Makerere University, 2012.
- J.R. Campell and J.F. Lasley, The science of Animals that serve humanity New York: Mc Graw Hill, 1975.

9. A. Clavijo, Y. Robinson, T. Booth and F. Munroe, Velogenic Newcastle in imported caged birds, 2000. Retrieved on 01/ 5/ 2012 from <http://www.kari.org/biennialconference/conference12/docs>
10. J.S. Chanta, A.O. Ademola, C.H. Emmanuel, R.A. Bello, P.O. Mbah, T.A. Adeniyi, F.K. Giscard, N. Emmanuel, F.Alain Le and P.M. Clauda, High Genetic Diversity of Newcastle Disease in poultry in West and Central Africa: Cocirculation of Genotype XIV and Newly Defined Genotype XVII and XVIII. *National institute of Health*. 51(7) 2250-2260, 2013.
10. M.K. Danda, D.M. Mwamachi, K. Leva and F. Jefa, Characterization of the Indigenous Chicken sub-sector in the Coastal Lowlands of Kenya, 2006. Retrieved on 01/5/ 2012 from <http://www.ncbi.nlm.nih.gov/pubmed/10816836>
11. M. Denscombe, *The good Research for small- scale Social Research Projects*. Buckingham: Open University Press, 1998.
12. L.K. Dontwi, F.T. Oduro, S.K. Amponsah and D. Appiah, Dynamics and control of viral diseases in poultry: A case Study of Ashanti Region in Ghana, 2011. Retrieved on 01/5/2012 from <http://hdl.handle.net/123456789/1242>
13. Food and Agricultural Organization, *The State of Food and Agriculture 2001*, Rome, Italy, 2001. Retrieved on 01/05/2012 from <http://www.fao.org/docrep/003/x9800e/x9800e14.htm>
14. M. Gay, *Education and Research Foundation Records*. North Carolina: Merdith College, 1987.
15. M.M. George, Epidemiology of Newcastle Disease and need to vaccinate local chickens in Uganda, 1992. In, P .B. Spradbrow (Ed). *Newcastle in Village chickens, control with thermo stable oral vaccine*, proceedings of the International workshop held in Kuala Lumpur, Malaysia 6-10 October 1991, Centre of International agricultural research (ACIAR).
16. C.M. Gichoni and J.G. Maina, *Poultry Production and Marketing*. Ministry of Livestock Production paper presented in Nairobi-Kenya, Nov. 23-27, 1992.
17. Government of Kenya, *Kenya Integrated Household Budgets survey (KIHBS) 2005/06.*, Basic report, Nairobi: Government Printer, Volume 1, 2007.
18. Government of Kenya, *Kisii South Development Plan 2008-2012*, Nairobi: Government printer, 2009.
19. Government of Kenya, *Ministry of Agriculture Farm Management Guidelines for Kisii South District*. Nairobi: Government printer, 2009.
20. E.F. Gueye, Village egg and meat production in Africa. *World's Poultry Science Journal* 54: 73-86, 1998.
21. E.F. Gueye, *Newcastle Disease in family poultry: Prospects for its control through ethnoveterinary medicine*, Dakar-Hann, Senegal, 2002.
22. C.D. Harvel, C.E. Mitchell, J.R. Waard, S. Altizer, A.P. Dobson, R.S. Ostfield and M.D. Samuel, Ecology-climate warming and disease risk for terrestrial and marine biota, 2002.
23. D.A. Higgins and K.F. Short ridge, *Newcastle disease in Tropical and developing countries.*. In D. J. Alexander (Ed), *Newcastle disease* Boston, Kluwer, pp.273-302, 1988
24. J.T. Houghton, Y. Ding, D.J. Griggs, M. Noguer, P.J. Van Der Linden, X. Dai, K. Maskell and C.A. Johnson, *Climate change: The scientific basis. Contribution of working group I to the third assessment report of the Intergovernmental Panel on Climate Change*, New York: Cambridge University Press, 2001.
26. Institute for International Cooperation in Animal Biologics, *Newcastle Disease*, College of Veterinary Medicine, Iowa State University, 2005.
27. Institute for International Cooperation in Animal Biologics, *Newcastle Disease. Avian paramyxovirus-1, Goose paramyxovirus infection*, College of Veterinary Medicine, Iowa State University, 2008.
28. Intergovernmental panel on climate change, IPCC, *Synthesis Report: Contribution of working groups I, II and III for the fourth assessment report of the Intergovernmental Panel on Climate Change*: Geneva, Switzerland, 2007.
29. H.A. Jensen and F. Dolberg, *A conceptual framework for using poultry as a tool in poverty alleviation. Livestock Research for Rural Development*, 2002. Retrieved on 07/13/2012, from <http://www.cipav.org.co/lrrd/lrrd15/5/jens155.htm>.
30. Johnston, Joe, and C. Robin, *Control of Newcastle Disease in Village Chickens with Oral V4 Vaccine*, Australia: Australian Centre for International Agricultural Research, 1991.
31. F.B. Kaingu, A.C. Kibor, R. Sharairo, H. Katima, J.O. Okeno, K. Wachenya, and A. K Kahi, Prevalence of gastro-intestinal Helminthes and Coccidian in indigenous chicken from different agro-climatic zones in Kenya, *African journal of Agricultural research*, 5 (6), pp458-462, 2010.
- Kenya National Bureau of Statistics, *2009 Kenya population census and Housing: Population distribution by administrative units*, Nairobi, 2009.
32. D.J. Kingston and D.C. Creswell, *Indigenous Chickens in Indonesia Population and Production Characteristics in Five Villages in West Java*, Winrock , International Arkansas USA, 1982.
33. M.A. Kingori, A.M. Wachira and J.K. Juitoek, *Indigenous Chicken Production in Kenya. A review. International journal of poultry* 9 (14) 309-316, 2010

34. C.A Kothari, Research Methodology: Methods and techniques (2 ed). New Delhi: Wishaw Prakashan, 1996.
35. B. Kouwenhoven, Newcastle disease virus infections in birds, 1993. Accessed on 17/10/2012 from <http://ci.vbi.edu/pathinfo/pathogens/NCDV>
36. J.E. Lancaster, Newcastle disease, a review of some of the literature published between 1926 and 1964, Ottawa: Canada, 1966.
37. H. M. Mai, O.D. Ogunsola and O.L. Obasi, Serological Survey of the Newcastle Disease and Infections Bursal Disease in Local Ducks and Guinea Fowls in Jos Plateau State, Nigeria. *Tropicaux journal* 57, 41-42, 2004.
38. P.A. Martin, The Epidemiology of Newcastle disease in village chicken's control with thermostable vaccines in proceedings, International workshop held in Kuala Lumpur, Malaysia. 1992.
39. J. McPeak, Confronting the risk of asset loss: What role do livestock transfers in northern Kenya play? *Journal of Development Studies*, 81, 415– 437, 2006.
40. Millar, Joanne, Photakoun and Viengxay, Livestock Development and Poverty Alleviation: Revolution or Evolution for Upland in Lao PRD? *Journal of Agricultural Sustainability* 6, 89-102, 2008.
41. Ministry of Agriculture, Farm Management Guidelines for Kisii South District year 2010-2011 Nairobi: Government printer, 2010.
42. Ministry of Livestock and Fisheries Development, MOLFD, Various Animal Reports. Nairobi: Government Printer, 2006.
43. Ministry of Livestock and Development, MOLD, Various Animal Reports, Nairobi: Government printer, 2007/2008.
44. J.C. Moreki, Family Poultry Production, poultry section, Department of Animal Production: Kenya , 2006.
45. W.B. Morgan and R.J.C. Munton, Agricultural Geography, New York: St Martin's Press 1972.
45. O.M. Mugenda and A.G. Mugenda, Research Methods: Quantitative and Qualitative approaches, Nairobi: Arc publishers, 2003.
46. G.Mukiibi-Muka, Epidemiology of Newcastle disease and the need to vaccinate local chickens in Uganda, 1992. In: Newcastle disease in village chickens (Spradbrow P.B. Editor), ACIAR proceedings No 39, Canberra, Australia.pp.155-158.
47. J.J. Musiime, The poultry industry in Kenya with particular reference to the Newcastle disease problem, 1992.. In, P.B. Spradbrow (Ed) Newcastle disease in village chickens, control with thermo stable oral Vaccines, Kuala Lumpur: Malaysia.
48. K. Mwiria and S.P. Wamahiu, Issues in Educational Research in Africa. Nairobi: E.A.E.P., 1995.
49. D.K. Nassiuma, Survey sampling: Theory and method, Nairobi University Press, Nairobi, 2000.
50. L.W. Njagi, P.N. Nyaga, P.G Mbuthia, I.C. Bebora, J N. Michieka, J.K. Kibe and U. M. Minga, Prevalence of Newcastle disease virus in village indigenous chickens in varied agro-ecological zones in Kenya. Department of Veterinary of pathology, microbiology, parasitology, Kangemi: Kenya, 2010.
51. A.E. Obayela and A. Andeniyi, The effect of climate on poultry in Ilorin Kwara State, Nigeria. University of Ibadan, Oyo state: Nigeria, 2006.
53. L.O. Okitoi, H.O. Ondwasy, D.N. Siamba, and D. Nkurumah, Traditional herbal preparations for indigenous poultry health management in Western Kenya, *Livestock Research for Rural Development* , 2007. Volume 19, Article #72. Retrieved June 30, 2010, from <http://www.lrrd.org/lrrd19/5/okit19072.htm>
53. H.O.K . Olabode, M.S. Damina, A.S. Ahmed, G.D . Moses and Y.S. Wungak, Retrospective Study of Newcastle Disease in Commercial Poultry Farms in Ilorin, Kwara State Nigeria: Department of Veterinary Microbiology, University of Abuja, Nigeria, 2012.
54. A.O. Olabode, A.G. Lamorde, N.N. Shidali and A.A. Chekwuedo, Village chickens and Newcastle disease in Nigeria, 1992. In: P. B. Spradbrow (Ed), Newcastle disease in village chickens. Control with Thermo stable oral vaccines proceedings, International workshop held in Kuala Lumpur: Malaysia
55. OIE, Newcastle disease. In: OIE manual of standards for diagnostic tests Vaccines 4th Ed., Paris: France, 2009.
56. E. C Okwor and D.C. EzeEpizootic Newcastle Disease in Local chickens reared in South East Savannah Zone in Nigeria. *International Journal of Poultry Science* 10(3), 212-215, 2011.
58. H. Ondwasy, H. Wesonga and I. Okitoi, Indigenous Chicken Production Manual. Nairobi: Kenya Research Institute, 1999.
59. B.D. Perry, T.F. Randolph, J. J. McDermott and P.K. Thornton, Investigating in animal health research to alleviate poverty, Nairobi: Kenya, 2002.
60. K.P. Riana, H.S. Susanne, G. E.Michael, D. Peter and E.F. Janet, Causal Inferences in disease ecology: Investigating ecological drivers of disease emergence, *International Journal of Poultry Science* Volume 6(8), 2008.
61. A. Sabuni, Climate change and parasitic and vector borne diseases, 27th October 2011. Retrieved on 3/5 2012 from <http://www.thepoultrysite.com>.

62. G. Shaw and D. Wheeler, Statistical techniques in geographical analysis, John and Wiley and Sons: New York, 1985.
 63. S.N. Seo and R. Mendelsohn, The impact of climate change on livestock management in Africa: A structural Ricardian analysis. CEEPA Discussion Paper No. 23. Centre for Environmental Economics and Policy in Africa, University of Pretoria, 2006.
 64. E.B. Sonaiya, Animal Science as a bridge between livestock farmers and consumers. Tropical Journal of Animal Science 2 (1):1-20, 1999a.
 65. E.B. Sonaiya, International network for family poultry development: Origins, activities, objectives and visions. In: Proc. Workshop on Poultry as a Tool in Poverty Eradication promotion of Gender Equality, 22-26 March. Landboskole, Denmark, 1999b. (also available at www.husdyr.kvl.dk/htm/php/tune99/4-sonaiya).
 66. Southern Africa Newcastle Disease Control Project SANDCP, Annual Plan 2005-06, GRM International: PYT LTD, 2005.
 67. P.B. Spradbrow, Epidemiology of Newcastle Disease and the Economics of its Control. In Dolberg, Frands and Poul Henning Petersen (ed). Poultry as a Tool in Poverty Eradication and Promotion of Gender Equality – Proceedings of a Workshop, 1999. Accessed on 07/07/2012 from
 68. D.E. Swayne and D.J. King, Avian Influenza and Newcastle Disease. J. A. Vet . Med. Association 222 (11) 1534-1540, 2003.
 69. H.T. Takakuwa, A. Takada, K. Okazaki and H. Kida, Potentially virulent Newcastle disease in migratory waterfowl populations. Japanese journal of veterinary research. 45(4), 207-215, 1998.
 70. O. Warner, Newcastle Disease (ND). In T. Blaha (Ed) Applied Veterinary Epidemiology, pp73-76. Elsevier: Amsterdam, 1989.
 71. E.G. White and F.T.W. Jordan, Newcastle disease, preventive medicine. Baillaire, Tindall & Cox ltd: London, 1963.
 72. B.A. Wilcox and R. Colwell, Emerging infectious Diseases. Biocomplexity as an inter disciplinary paradigm. Ecohealth vol. 2(4) pp.244-257, 2005.
- World Health Organisation, W.H.O, Using Climate to Predict Infectious Disease Outbreaks: A Review. Geneva, 2004.