



DSPACE

<https://dspace.org/>

Incidence of pests and diseases in tree nurseries and plantations in Kimondi forest, Nandi county, Kenya

Ondieki, Joseph A; Sirmah, Peter; Hitimana, Joseph

2021-03-05

East African Journal of Forestry and Agroforestry

<http://ir-library.kabianga.ac.ke/handle/123456789/326>

Ondieki, J. A., Sirmah, P., & Hitimana, J. (2021). Incidence of Pests and Diseases in Tree Nurseries and Plantations in Kimondi Forest, Nandi County, Kenya. East African Journal of Forestry and Agroforestry, 3(1), 18-28.



East African Journal of Forestry & Agroforestry

eajfa.eanso.org

Volume 3, Issue 1, 2021

Print ISSN: 2707-4315 | Online ISSN: 2707-4323

Title DOI: <https://doi.org/10.37284/2707-4323>



EAST AFRICAN
NATURE &
SCIENCE
ORGANIZATION

Original Article

Incidence of Pests and Diseases in Tree Nurseries and Plantations in Kimondi Forest, Nandi County, Kenya

Joseph Atanga Ondieki^{1*}, Dr. Peter Sirmah (Ph.D)², & Dr. Joseph Hitimana (Ph.D)³

¹ Department of Agroforestry, Environmental Studies and Integrated Natural Resource Management, University of Kabianga, P. O. Box 2030-20200, Kericho, Kenya.

* ORCID: <https://orcid.org/0000-0002-6619-5870>; Author for Correspondence email: psirmah@kabianga.ac.ke.

Article DOI: <https://doi.org/10.37284/eajfa.3.1.287>

Date Published: ABSTRACT

05 March 2021

Keywords:

Pests,
diseases,
tree nurseries,
plantations,
Kimondi forest,
Kenya.

Increasing demand for wood is putting pressure on forest resources that are equally under threat from insect pests and diseases. Reported average annual forest loss stands at 0.2% globally, 0.8% in Africa, and 1.6% in Kenya. To meet the increasing demand for forest products in Kenya, the government and private sector have established plantations of non-native tree species dominated by *Eucalyptus species*, *Cupressus lusitanica*, and *Pinus patula*. To ensure successful forest plantation establishment and management schemes, there is a need for sufficient knowledge and understanding of tree growth conditions and threats including pests and diseases. This study aimed at determining the incidence of plantation and tree nurseries in Kimondi Forest, Kenya. Tree plantations were mapped into 2.5 Ha rectangular portion transects parallel to the forest roads. In randomly selected portions, observations were carried out for disease and pest signs and symptoms on various tree parts (leaves, stem, roots, fruits, and twigs). In the tree nurseries, 3 m × 1 m rectangular quadrants were placed on seedling beds and similar observations were made. Collected data indicated a high incidence of nursery seedling pests (5.3 % leaves and 5.1% stems) on *Eucalyptus species* and least on *P. patula* (2.4% leaves and 3% stems). Higher incidence of plantation pests (35.0% leaves of *Eucalyptus sps.*) and least on *P. patula* stems (1.2%) were recorded. On the other hand, twelve (12.0%) of *C. lusitanica* and (1.8%) *Eucalyptus species* plantation twigs were infested by pests. A high incidence of nursery seedling disease (9.8%) was observed on *C. lusitanica* leaves and least on *P. patula* stems (3.6%). Higher incidence of plantation diseases (32 %) on leaves of *Eucalyptus sps.* and least on *C. lusitanica* stem (1.4%) was recorded. In both tree nurseries and plantations, roots and fruits remained free from pests and disease. Major tree

pests and diseases identified in Kimondi forests include (Human, wildlife, livestock, *Cinara cupressi*, *Gonipterus scutellatus*, *Pinus pini*, and *Leptocybe invasa*) and (damping-off, *Fusarium wilt*, *Botryosphaeria* canker, cypress canker, and *Mycosphaerella* spp.) respectively. These results suggest a need for regular monitoring and intervention measures to control pest and disease infestation in the Kimondi forest.

APA CITATION

Ondieki, J. A., Sirmah, P., Hitimana, J. (2021). Incidence of Pests and Diseases in Tree Nurseries and Plantations in Kimondi Forest, Nandi County, Kenya *East African Journal of Forestry and Agroforestry*, 3(1), 18-28. <https://doi.org/10.37284/eajfa.3.1.287>

CHICAGO CITATION

Ondieki, Joseph Atanga, Peter Sirmah, Joseph Hitimana. 2021. "Incidence of Pests and Diseases in Tree Nurseries and Plantations in Kimondi Forest, Nandi County, Kenya". *East African Journal of Forestry and Agroforestry* 3 (1), 18-28. <https://doi.org/10.37284/eajfa.3.1.287>.

HARVARD CITATION

Ondieki, J. A., Sirmah, P., Hitimana, J. (2021) "Incidence of Pests and Diseases in Tree Nurseries and Plantations in Kimondi Forest, Nandi County, Kenya", *East African Journal of Forestry and Agroforestry*, 3(1), pp. 18-28. doi: 10.37284/eajfa.3.1.287.

IEEE CITATION

J. A. Ondieki, P. Sirmah, J. Hitimana, , "Incidence of Pests and Diseases in Tree Nurseries and Plantations in Kimondi Forest, Nandi County , Kenya", *EJFA*, vol. 3, no. 1, pp. 18-28, Mar. 2021.

MLA CITATION

Ondieki, Joseph Atanga, Peter Sirmah, Joseph Hitimana. "Incidence of Pests and Diseases in Tree Nurseries and Plantations in Kimondi Forest, Nandi County, Kenya". *East African Journal of Forestry and Agroforestry*, Vol. 3, no. 1, Mar. 2021, pp. 18-28, doi:10.37284/eajfa.3.1.287.

INTRODUCTION

Increasing demand for wood and wood products has exerted pressure on natural forest resources throughout the world (FAO, 2007). These resources are currently under threat due to other factors including clearing forest land for agricultural production or settlements. According to FAO (2007), the world average annual forest losses stand at 0.2%, while that of Africa is about 0.8%.

The need to accommodate the increasing demand for tree products such as fibre and wood in Africa has therefore led to an increasing establishment of forest plantations using non-native tree species such as *Eucalyptus*, *Cupressus lusitanica* and *Pinus patula* species (Evans, 1992). These plantations have an important economic and social bearing for several countries in Africa providing a source of employment, wealth creation, the production of export capital as well as fuel and construction timber (FAO, 2007). Despite their importance, forest plantations using non-native species (exotic)

are fragile and face unique threats mostly from pests and disease (Wingfield *et al.*, 2001; Cock, 2003) due to a limited genetic base (Wingfield *et al.*, 2008). Documented examples of major disease outbreaks in the history of forest plantations include *Dothistroma* needle blight of *Pinus* species in Chile, New Zealand and Kenya (Gibson, 1975); *Phytophthora pinifolia* in Chile (Durán *et al.*, 2008); conifer aphid (*Cinara cupressivora*) on *Cupressus* species in Eastern and Central Africa (Murphy, 1998; Day *et al.*, 2003); the sirex wood wasp (*Sirex noctilio*) on *Pinus* species in South Africa (Tribe, 2003; Hurley *et al.*, 2007), and the blue gum chalcid (*Leptocybe invasa*) on *Eucalyptus* species in many countries in Africa, Asia, the Middle East and Europe (Mutitu, 2003; Mendel *et al.*, 2004; Nyeko, 2005; EPPO, 2006). *Dothistroma septosporum* led to the termination of *P. radiata* planting in many African countries, particularly those in East Africa (Gibson *et al.*, 1964; Barnes *et al.*, 2004). Similarly, the susceptibility of *Eucalyptus* to the snout beetle, *Gonipterus scutellatus* and *Mycosphaerella* leaf disease led to the partially discontinued planting of

this species in South Africa (Lundquist et al., 1987). The afforestation examples illustrate how forest pests and diseases raise serious concerns to developers of tropical tree plantation enterprises. Unfortunately, FAO observed that many people working with trees do not pay systematic attention to tree health until trees have died and it is too late for intervention (FAO, 2003).

Outbreaks of serious pests and diseases may still be a major threat to future plantation development in Kenya. Example devastating outbreaks of Psyllid, *Heteropsylla cubana* in plantations of *Leucaena leucocephala* in South East Asia, *conifer aphids* in plantations in East Africa and *eucalyptus* leaf diseases are only recent history (Nair, 2000).

In order to contain pest and pathogen problems, there is a need for knowledge and information regarding the occurrence of nursery and plantation insect pests and diseases. This study aimed at


evaluating the incidence of pests and diseases of tree nurseries and plantations of Kimondi Forest, Nandi County, Kenya.

RESEARCH METHODOLOGY

Study Area

This study was carried out in the Kimondi forest located in Nandi County, Kenya (Figure 1). It lies on latitude $0^{\circ} 18'$ and $0^{\circ} 32'$ N and longitude $37^{\circ} 05'$ to $37^{\circ} 23'$ E in the North Rift Conservancy. The soils are characterized as very shallow to deep and dark yellowish-brown to very dark brown, respectively. Soil types range from friable sandy loam to sandy clay loam and friable gravelly sandy clay loam to sand clay with 5 – 50% fine to medium iron and manganese concentrations (Otieno et al., 2014)).

Figure 1: Map of the Study Area



Source: Adopted from KFS.

The Kimondi forest is mainly drained by both Kimondi and Sirua rivers which merge further downstream to form River Yala that finally flows into Lake Victoria (Mitchell, 2004). The forest is the source of water for the recently inaugurated Nzoia cluster II water project (supplies water to Kakamega town and its surroundings) and the Kobujoi water supply project (supplies water to Kobujoi and its surrounding) (Simon *et al.*, 2016). Other rivers that drain water out of the Kimondi forest include Mokong, Orobo, Kundos, Gorgor, Chemogonja and Cheptaburur. These rivers are perennial and therefore provide water for domestic and industrial use by the adjacent forest communities as well as have waterfalls, which can be harnessed for hydroelectric power generation (District Development Plan, Nandi South 2008-2012, GoK, 2008).

Target Population

The study targeted pests and diseases in both tree nursery seedlings and plantation species of *Cupressus lusitanica* (678.3 Ha), *Pinus patula* (147.5 Ha) and *Eucalyptus species* (48.8 Ha).


Sampling in Tree Nurseries

Tree seedlings within 10 portions of size 3 m × 1 m were sampled for this study according to Sokal and Rohlf (2012) as follows: Nursery beds of the three tree species (*C. lusitanica*, *P. patula*, and *Eucalyptus species*) were separately divided into equal portions of 3 m by 1 m along the nursery lengths. Each portion was numbered; then, 10 portions were randomly selected for the purpose of this study. Incidence of nursery pests and diseases were observed on the roots, leaves and stems of the seedlings within each portion.

Sampling in the Forest Plantations

The targeted plantations (*Eucalyptus*, *C. lusitanica* and *P. patula*) were separately mapped into 2.5 Ha rectangular plots (Figure 2). The plots were positioned in such a way that its long axis ran parallel with the planting rows.

Figure 2: Configuration of the 2.5 Ha Sample Plot Within a Plantation



The ratio of the length of road within the compartment (L) to the compartment area (A) was the same as the length of road (Ls) within the 2.5 Ha sample area (As).

Ground line transect walk was carried out along the road and a take-off point was randomly chosen along the distance abutting each of the 2.5 Ha sample plots per plantation species. Observations were made on 20 consecutive trees for incidences of pests and disease incursion. Across several rows, a further 20 consecutive trees were inspected for damage on the return journey to the road.

Determination of Diseases and Pests Incidence

In each sampled portion (nursery beds or plantations) number of diseased and pest-infested roots, leaves, fruits, twigs and stems were recorded separately and expressed as the proportion of total number in each set respectively according to Anonymous (2006) as follows:

$$\text{Leaf incidences(\%)} = \frac{\text{Number of infected leaves}}{\text{Total leaves observed in a portion}} \times 100 \dots \dots \dots [1]$$

$$\text{Fruit incidence (\%)} = \frac{\text{Number of infected fruits}}{\text{Total fruits observed in a portion}} \times 100 \dots \dots \dots [2]$$

$$\text{Twigs incidences(\%)} = \frac{\text{Number of infected twigs}}{\text{Total twigs observed in a portion}} \times 100 \dots\dots\dots [3]$$

$$\text{Root incidences(\%)} = \frac{\text{Number of infected leaves}}{\text{Total leaves observed in a portion}} \times 100 \dots\dots\dots [4]$$

$$\text{Stem incidences(\%)} = \frac{\text{Number of infected leaves}}{\text{Total leaves observed in a portion}} \times 100 \dots\dots\dots [5]$$

Photographs of pest and disease-infested plant parts were taken and used for identification purposes.

RESULTS AND DISCUSSION

Incidence of Tree Nursery and Plantation Pests

Table 1 reports the incidence of nursery pests in Kimondi forest tree nurseries. Five (5.3 %) of *Eucalyptus* leaves were damaged by pests in comparison with 4.7% in *C. lusitanica* and 2.4 % in

P. patula. Roots for all the seedlings were free from pests.

Table 1: Incidence of Tree Nursery Pests

Species	(%) Incidence of Pests		
	Leaves	Stems	Roots
<i>P. patula</i>	2.4	3.0	0.0
<i>C. lusitanica</i>	4.7	5.0	0.0
<i>Eucalyptus</i>	5.3	5.2	0.0

Table 2 reports the incidence of pests in the Kimondi forest plantation. Thirty-Five per cent (35%) of *Eucalyptus* leaves were damaged by pests in comparison with 11.5% in *C. lusitanica* and 9.6 % in *P. patula*. Twelve per cent (12%) of *C. lusitanica* twigs were damaged by pests in comparison to a lower margin (1.8%) observed in *Eucalyptus* species. Fruits for all the tree species were free from pests.

Table 2: Incidence of Forest Plantation Pests

Species	% Incidence of Pests			
	Leaves	Stem	Twigs	Fruits
<i>P. patula</i>	9.6	1.2	8.1	0.0
<i>C. lusitanica</i>	11.5	1.3	12.0	0.0
<i>Eucalyptus species</i>	35.0	5.6	1.8	0.0

Minor pest attack on *Eucalyptus* (5.6%), *P. patula* (1.3%) and *C. lusitanica* (1.2%) stems were observed. These results suggest that leaves of both the tree nurseries and plantation species are mainly targeted by pests in comparison to the stems, fruits and roots. Such observation has been reported in other similar studies (Nsolomo & Venn, 1984; Nyeko & Nakabonge, 2008; Wingfield *et al.*, 2010).

Incidence of Tree Nursery and Plantation Diseases

Table 3 reports the incidence of diseases in Kimondi forest tree nurseries. 9.8% of *C. lusitanica* leaves were damaged by diseases in comparison with 8.0% in *P. patula* and 5.6% in *Eucalyptus* tree species.

Table 3: Incidence of Tree Nursery Diseases

Species	(%) Incidence of Diseases		
	Leaves	Stems	Roots
<i>P. patula</i>	8.0	5.1	0.0
<i>C. lusitanica</i>	9.8	3.6	0.0
<i>Eucalyptus species</i>	5.6	7.7	0.0

Based on Table 3, 7.7% of *Eucalyptus species* stems were damaged by diseases in comparison to 3.6% observed in *C. lusitanica* and 5.1% in *P. patula*. Roots of all tree seedling species remained free from

diseases. Such observations have been reported in other similar studies (Gichora *et al.*, 2017).

Table 4 reports the incidence of diseases in the Kimondi forest plantation. Thirty-Five (32%) of *Eucalyptus* leaves were damaged by pests in comparison with 12% in *C. lusitanica* and 21 % in

P. patula. Fourteen (14.6%) of *C. lusitanica* twigs were damaged by pests in comparison to a higher margin (17%) observed in *Eucalyptus* species. Fruits for all the tree species were free from pests, while minimal damage on stems (8.5%) on *Eucalyptus* species, (1.4%) on *C. lusitanica* and 3.2% on *P. patula* was observed.

Table 4: Incidence of Forest Plantation Diseases

Species	(%) Incidences of Diseases			
	Leaves	Stem	Twigs	Fruits
<i>P. patula</i>	21.3	3.2	7.8	0.0
<i>C. lusitanica</i>	12.0	1.4	17.7	0.0
<i>Eucalyptus species</i>	32.0	8.5	14.6	0.0

Major Pests and Diseases in Kimondi Forest

Table 5 reports the major pests identified in the Kimondi forest. *Leptocybe invasa* was observed on young seedlings of *Eucalyptus species* in the

nursery (Plate 1a & b) as small circular galls on the midribs of the leaves. *Leptocybe invasa* lays eggs on the bark of shoots or the midribs of leaves; eggs develop into the minute, white, legless larvae within the host plant.

Table 5: Major Pests Identified in Kimondi Forest

Pest	Observed Symptoms	Affected Tree Species
<i>Leptocybe invasa</i>	Pinhole on leaves, galls on mid-ribs and petioles	<i>Eucalyptus</i> leaves
Termites	Nearby termite mounds, stems debarked at the base of the plant	All species
Livestock/wildlife	De-budding, cut stems and branches, debarking, browsings on leaves	All species
<i>Pinus pini</i>	Defoliation, dieback, needle shading, dead trunks	<i>Pinus patula</i>
<i>Goniapterus scutellatus</i>	Leaf defoliation, scalloped leaf edges, dieback of shoot tips	<i>Eucalyptus</i>
<i>Cinera cupressi</i>	Galls	<i>Cupressus lusitanica</i>
Human	Logging, debarking, cuttings, uprooting	All species

Plate 1: a) Galls of *Leptocybe invasa* on Eucalyptus Leaves; b) Debarked *P. Patula* Stem**a****b**

Damage to trees is caused when the developing larvae of *Leptocybe invasa* produce galls on the leaf midribs, petioles and twigs. The galls cause the twigs to split, destroying the cambium. Small circular holes indicating exit points of adults from pupae are common on the galls. Repeated attacks lead to loss of growth and vigour in susceptible trees. Severely attacked seedlings show gnarled appearance, stunted growth, lodging, dieback and eventually die (Mendel *et al.*, 2004; Nyeko & Nakabonge, 2008; Jhala, Patel & Vaghela, 2010; Petro, 2015; de Souza *et al.*, (2014).

Termites attack on both *Eucalyptus species* and *C. lusitanica* seedlings was associated with debarked shoots slightly above and below the base level and

the presence of anti-hills around the tree nursery. Petro (2015) previously observed termites as a major pest of nursery seedlings and young plantation of *Eucalyptus species* below age five. Termites debark the stems below and slightly above ground level, thus affecting the functioning of the physiological process of plant structures, thus leading to the death of seedlings in the nursery (Nyeko & Nakabonge, 2008). Other pests included humans, sap sacking *Cinera cupressi* and leaf defoliation caused by *Pinus pini*. These results are in agreement with the findings of Nyeko and Nakabonge (2008) and Gichora *et al.* (2017). Major diseases in the Kimondi forest were identified by observing disease signs and symptoms. Results are presented in Table 6.

Table 6: Major Diseases Identified in Kimondi Forest

Disease	Observed Symptoms	Affected Tree Species
Damping-off	Wilting, rotting stems, toppling or drying of stems	Young seedlings of all species
<i>Fusarium wilt</i>	Chlorosis of terminal needles wilted flaccid needles	<i>Pinus patula</i>
<i>Botryosphaeria canker</i>	Kino cracks, resin exudation, dieback on tips, coppice failure, fruiting bodies	<i>Eucalyptus</i>
<i>Cypress canker</i>	Fruiting bodies, yellowing of needles, dieback, oozing resin, stem lesion	<i>Cupressus lusitanica</i>
<i>Mycosphaerella sps</i>	Pale brown leaf margins	<i>Eucalyptus</i>

Damping-off disease associated with wilting and rotting stems was prevalent in young seedlings (Plate 2b). However, sometimes the yellowing and wilting of pine seedlings in some nurseries are associated with delayed pricking out, overstay in the nursery or use of insufficient mycorrhizal soil

(Drenth & Sendall, 2001), but this was never the case. *Eucalyptus* seedlings in the nursery as well as in the plantation displayed dieback and stem canker symptoms associated with *Botryosphaeria* infection (Plate 2a).

Plate 2: a) Cracks, Kino and Resin Exudation on Eucalyptus Stem; b) Wilting of *C. lusitanica* Seedlings due to Damping Off Disease



a



b

Botryosphaeria infection was characterized by dieback of the growing tips, coppice failure and stem cankers. Old trees showed extensive kino accumulation, wood rot and fruiting bodies of *Botryosphaeria* spp. Cypress canker characterized by fruiting bodies on infected trees, yellowing of

needles, and dieback of branches, oozing resin, stem lesion and trunks were observed on *C. lusitanica* plantation. *Mycosphaerella* spp characterized by irregular pale, brown colouration on leaves with distinct red-brown leaf margin was observed (Plate 3b).

Plate 3: a) Stem Canker of *C. lusitanica*; b) *Mycosphaerella* Sps on *Eucalyptus* Leaves**a****b**

CONCLUSION

A high incidence of nursery seedling pests (5.3% leaves and 5.1% stems) was observed on *Eucalyptus* species and least on *P. patula* (2.4% leaves and 3% stems). Higher incidence of plantation pests (35.0% leaves of *Eucalyptus* spp.) and least on *P. patula* stems (1.2%) were recorded. On the other hand, twelve (12.0%) of *C. lusitanica* and (1.8%) *Eucalyptus* species plantation twigs were infested by pests. A high incidence of nursery seedling disease (9.8%) was observed on *C. lusitanica* leaves and least on *P. patula* stems (3.6%). Higher incidence of plantation diseases (32 %) leaves of *Eucalyptus* spp. and least on *C. lusitanica* stem (1.4%) was recorded. In both tree nurseries and plantations, roots and fruits remained free from pests and disease, respectively. Major tree pests and diseases identified in Kimondi forests include (Human, wildlife, livestock, *Cinera cupressi*, *Gonipterus scutellatus*, *Pinus pini* and *Leptocybe invasa*) and damping off, *Fusarium* wilt, *Botryosphaeria* canker, cypress canker and *Mycosphaerella* spp.) respectively. These results suggest a need for intervention measures to control pest and disease infestation in the Kimondi forest.

REFERENCE

- Anonymous. (2006). Survey methodology for Bacterial blight of pomegranate, NCRP publication, pp.14
- Barnes, I., Crous, P. W., Wingfield, B. D., & Wingfield, M. J. (2004). Multigene phylogenies reveal that red band needle blight of *Pinus* is caused by two distinct species of *Dothistroma*, *D. septosporum* and *D. pini*. *Studies in Mycology*, 50(2), 551-565.
- Cock, M. J. W. (2003). *Biosecurity and Forests: An introduction with particular emphasis on forest pests*. Forest Health and Biosecurity Working paper FBS/2E. Rome, Italy: FAO.
- Day, R. K., Kairo, M. T. K., Abraham, Y. J., Kfir, R., Murphy, S. T., & Mutitu, K. E. (2003). Biological control of Homopteran pests of conifers in Africa. In Neuenschwander, P. Borgemeister, C. & Langewald, J. (Eds), *Biological control in IPM systems in Africa* (pp. 101-112). CAB International, Wallingford, UK.
- de Souza, M. D., Rezende, E. H., Alves, T., Bandeira, R. R., & Fernandes, A. D. M. (2014). First report of *Leptocybe invasa* Fisher and

- LaSalle (Hymenoptera: Eulophidae) in Mozambique. *African Journal of Agricultural Research*, 9(49), 3555-3558.
- Drenth, A., & Sendall, B. (2001). Practical guide to detection and identification of Phytophthora. *Tropical Plant Protection*, 1, 32-33.
- Durán, A., Gryzenhout, M., Slippers, B., Ahumada, R., Rotella, A., Flores, F., ... & Wingfield, M. J. (2008). Phytophthora pinifolia sp. nov. associated with a serious needle disease of Pinus radiata in Chile. *Plant Pathology*, 57(4), 715-727.
- European and Mediterranean Plant Protection Organization (EPPO). (2006). *EPPO Alert List: Leptocybe invasa (Hymenoptera: Eulophidae) – Blue Gum chalcid*. Retrieved on 26th March 2018 from http://www.eppo.org/QUARANTINE/Alert_List/insects/Leptocybe_invasa.htm.
- Evans, J. (1992). *Plantation forestry in the tropics: tree planting for industrial, social, environmental, and agroforestry purposes*. Oxford University Press.
- FAO. (2003). *An illustrated guide to the state of health of trees Recognition and interpretation of symptoms and damage*. Diagnostic and Advisory Service CABI Bioscience Egham, Surrey, United Kingdom
- FAO. (2006). *Global Forest Resources Assessment 2005 – progress towards sustainable forest management*. Forestry Paper No. 147. Rome, Italy. Available at: <http://www.fao.org/docrep/008/a0400e/a0400e00.htm>
- FAO. (2007). Forest Pest Species Profile: *Heterosyllacubana*, Crawford, 1914. Rome: FAO. www.fao.org/forestry/13564-1-0.pdf
- Gibson, I. A. S., Christensen, P. S., & Munga, F. M. (1964). First observations in Kenya of a foliage disease of pines caused by Dothistroma pini Hulbary. *The Commonwealth Forestry Review*, 31-48.
- Gibson, I.A.S. (1975). *Diseases of forest trees widely planted as exotics in the tropics and southern hemisphere. Part 1, important members of the Myrtaceae, Leguminosae, Verbinaceae and Meliaceae*. Commonwealth Mycological Institute and Forestry Institute. Kew and Oxford.
- Gichora, M., Kojwang, H., & Bosu, P. (2017). *The status and trends of forest and tree pests and diseases management in Africa*. African Forest Forum Working Papers (3)3. African Forest Forum, Nairobi.
- GOK. (2008). *Kenya Forest Service Board Paper: Revision of royalties for financial year 2008/09*. Nairobi, KE: KFS.
- Hurley, B. P., Slippers, B., & Wingfield, M. J. (2007). A comparison of control results for the alien invasive woodwasp, Sirex noctilio, in the southern hemisphere. *Agriculture Forest Entomology*, 9, 159-171.
- Jhala, R. C., Patel, M. G., & Vaghela, N. M. (2010). Effectiveness of insecticides against blue gum chalcid, Leptocybe invasa Fisher & La Salle (Hymenoptera: Eulophidae), infesting eucalyptus seedlings in middle Gujarat, India. *Karnataka Journal of Agricultural Sciences*, 23(1), 84-86.
- Lundquist, J. E., Purnell, R. C., & de Wet, D. R. (1987). Effects of Mycosphaerella leaf spot on the growth of Eucalyptus nitens. *Plant Disease*, 71(11), 1025-1029.
- Mendel, Z., Protasov, A., Fisher, N., & La Salle, J. (2004). Taxonomy and biology of Leptocybe invasa gen. & sp. n.(Hymenoptera: Eulophidae), an invasive gall inducer on Eucalyptus. *Australian Journal of Entomology*, 43(2), 101-113.
- Mitchell, N. (2004). *Exploitation and disturbance history of Kakamega Forest, western Kenya*. BIOTA East Report number 1
- Murphy, S. T. (1998). Protecting Africa's trees. *UNASYLVA-FAO*, 57-61.

- Mutitu, K. E. (2003). *A pest threat to Eucalyptus species in Kenya*. KEFRI Technical Report. Nairobi, KEFRI.
- Nair, K. S. S. (ed.) (2000). *Insect Pests and Diseases in Indonesian Forests: An Assessment of the Major Threats, Research Efforts and Literature*. Bogor, Indonesia: Center for International Forestry Research.
- Nsolomo, V. R., & Venn, K. A. R. E. (1994). Forest fungal diseases of Tanzania: background and current status. *Norwegian Journal of Agricultural Sciences*, 8, 189-189.
- Nyeko, P. & Nakabonge, G. (2008). *Occurrence of pests and diseases in tree Nurseries and plantations in Uganda*. Kampala, Uganda: the Sawlog Production Grant Scheme (SPGS)
- Nyeko, P. (2005). The cause, incidence and severity of a new gall damage on Eucalyptus species at Oruchinga refugee settlement in Mbarara district, Uganda. *Uganda Journal of Agricultural Sciences*, 11, 47-50.
- Oballa, P. O., Konuche, P. K. A., Muchiri, M. N. and Kigomo, B. N. (2010). *Facts on growing and use of Eucalyptus in Kenya*. KEFRI, Nairobi, Kenya.
- Otieno, N. E., Sajita, N., & Shitandayi, D. (2014). Response of a globally endangered canopy insectivore to habitat degradation in an East African tropical rainforest: The role of differential forest protection levels. *International Journal of Biodiversity and Conservation*, 6(3), 290-300.
- Petro, R. (2015). *Effects of eucalyptus gall wasp, leptocybe invasa (Hymenoptera: eulophidae) on growth and wood basic density of some eucalyptus species, Tanzania*. Doctoral Thesis, Sokoine University of Agriculture.
- Simon, R. K., Kosgei, A. J., & Kireger, E. (2016). Economic valuation of underutilized forestry Products in South Nandi Forest. *International Journal of Research in Applied, Natural and Social Sciences*. 4(8), 29-36.
- Sokal, R. R., & Rohlf, F. J. (2012). *Biometry*, 4th ed. New York. NY: WH Freeman and Company.
- Tribe, G. D. (2003). Biological control of defoliating, and phloem- or wood-feeding insects in commercial forestry in southern Africa. . In Neuenschwander, P. Borgemeister, C. & Langewald, J. (Eds), *Biological control in IPM systems in Africa* (pp. 113-129). CAB International, Wallingford, UK.
- Wingfield, M. J., Slippers, B., Hurley, B. P., Coutinho, T. A., Wingfield, B. D., & Roux, J. (2008). Eucalypt pests and diseases: growing threats to plantation productivity. *Southern Forests: a Journal of Forest Science*, 70(2), 139-144.
- Wingfield, M.J., Slippers, B., & Wingfield, B.D. (2010). Novel associations between pathogens, insects and tree species threaten world forests. *New Zealand Journal of Forest Sciences*, 40, S95-S103.
- Wingfield, M. J., Slippers, B., Roux, J., & Wingfield, B. D. (2001). Worldwide movement of exotic forest fungi, especially in the tropics and the southern hemisphere: This article examines the impact of fungal pathogens introduced in plantation forestry. *Bioscience*, 51(2), 134-140.